

# American Journal of Science

DECEMBER 1985

## GEOCHEMISTRY OF LATERITES, STABILITY OF Al-GOETHITE, Al-HEMATITE, AND $\text{Fe}^{3+}$ -KAOLINITE IN BAUXITES AND FERRICRETES: AN APPROACH TO THE MECHANISM OF CONCRETION FORMATION

YVES TARDY\* and DANIEL NAHON\*\*

**ABSTRACT.** In most lateritic weathering profiles, crystallized iron-rich oxihydroxides are found to be solid solutions of goethite-diaspore or hematite-corundum. Moreover, kaolin minerals are generally found to contain small quantities of structural iron and can also be considered as solid solutions of  $\text{Fe}^{3+}$  and  $\text{Al}^{3+}$ -kaolinite end-members. In bauxites, ferricretes, and in the top of lateritic profiles, important nodule formation takes place. The nodule formation taking place within a kaolinite soil-matrix is always made up of Al-hematite which replaces the clay through an epigenetic relation. From nodules to pisolites, the centripetal evolution results in the formation of Al-goethite at the expense of earlier formed hematite.

The distribution of  $\text{Fe}^{3+}$ -kaolinite, Al-goethite, and Al-hematite as well as the iron and aluminum contents is controlled by silica activity, pore size distribution, and the activity of water. A model calculation, based on solid solution equilibria and minimization of the Gibbs free energy in the system  $\text{Fe}_2\text{O}_3$ - $\text{Al}_2\text{O}_3$ - $\text{SiO}_2$ - $\text{H}_2\text{O}$ , is proposed. The model calculates the amounts of coexisting phases and the distribution of Al and Fe in goethite, hematite, and kaolinite. It is shown that when kaolinite controls the system, the amounts of aluminum in goethite and hematite increase when the activity of silica decreases. Also, when the water activity diminishes, the Al-content in goethite increases. For low water activities, hematite replaces goethite, while boehmite replaces gibbsite. Equilibrium conditions and element distributions are also shown to be dependent on the amounts of available water in the system so that results of calculations are compatible with natural observations.

### INTRODUCTION

Under warm, humid, or semi-humid conditions such as those prevailing under tropical or mediterranean climates, red and yellow are the dominant kaolinitic soil colors. In contrast, soils of temperate regions are dominantly yellowish colored. The high chroma-colors of the lateritic soils are determined by the amounts of ferrihydrite, hematite, and goethite. Goethite is by far the most widespread form of pedogenic iron oxide and

\* Institut de Géologie et Centre de Sédimentologie et de Géochimie de la Surface, 1, rue Blessig, 67084 Strasbourg Cedex, France

\*\* Laboratoire de Pétrologie de la Surface, U.A. 721 du C.N.R.S., 40, avenue du Recteur Pineau 86022, Poitiers Cedex, France

may originate and persist in widely diverse environments (Schwertmann, Fischer, and Taylor, 1974; Schwertmann and Taylor, 1977). The zonal and intrazonal occurrence of goethite and hematite has been documented in many places (Taylor and Graley, 1967; Schwertmann, 1969; Fölster, Noshrefi, and Ojenuga, 1971; Lamouroux, 1972; Davey, Russel, and Wilson, 1975; Nahon, 1976; Kämpf and Schwertmann, 1983).

Since Correns and Von Engelhardt (1941), Norrish and Taylor (1961), and Bardossy (1968), substitutions of Al for Fe in the structure of goethite and hematite have been found in bauxites, ferricretes, laterites, and in most of the natural profiles in which these minerals form. Correlatively, since Boesman and Schoemacker (1961) many natural kaolinites have been determined as  $\text{Fe}^{3+}$ -substituted. Amounts of iron in kaolinites are generally very small but highly dependent on the physicochemical properties of these clay minerals (Yvon and others, 1981; Cases and others, 1981).

This paper is an attempt to review the distribution, range of composition, and conditions of formation of Al-goethites, Al-hematites, and  $\text{Fe}^{3+}$ -kaolinites in natural tropical weathering profiles, including bauxites and ferricretes.

#### SUBSTITUTION RANGES AND METHOD OF DETERMINATION

*A. Determination of the substitution of  $\text{Al}^{3+}$  for  $\text{Fe}^{3+}$  in goethites and hematites.*—Measurements of the degree of substitution of  $\text{Al}^{3+}$  for  $\text{Fe}^{3+}$  in goethite and hematite have been performed by Mössbauer spectrometry since Janot and Gibert, 1970. Another method is based on the relation between shifts of X-ray diffraction peaks and degrees of substitution according to the Vegart Law (Norrish and Taylor, 1961; Thiel, 1963). Deviations from the Vegart Law have been found by Perinet and Lafont (1972a, b) and verified by Schwertmann, Fitzpatrick, and Le Roux (1977) for hematites. Parameter  $\Delta a$  can be calculated from the X-ray diffractogram, and its deviation  $\Delta a(0.001 \text{ \AA})$  is a function of the mole fraction of  $\text{Al}_2\text{O}_3$  within hematite (X percent) according to the relation:

$$\Delta a = 0.022 X^2 + 1.82 X$$

*B. Determination of the substitution of  $\text{Fe}^{3+}$  for  $\text{Al}^{3+}$  in kaolinites.*—The substitution of  $\text{Fe}^{3+}$  into kaolinite has been studied by Mössbauer spectrometry (Malden and Meads, 1967; Janot, Gibert, and Tobias, 1973). It seems also that quite precise results may be obtained by Electron Spin Resonance (ESR), elsewhere called EPR or RPE. An ESR spectra of  $\text{Fe}^{3+}$ -kaolinite shows three types of signals among which two are significantly used. A first group of resonance signals is located around  $g = 4.67$  Gauss. This group includes two main peaks called internal ( $S_I$ ) and external ( $S_E$ ). The amount of  $\text{Fe}^{3+}$  substituted is proportional to the sum of the areas of these two peaks ( $S_I + S_E$ ). The kaolinite cristallinity is inversely correlated with the area of the internal peak ( $S_I$ ) and positively correlated with the area of the external peak ( $S_E$ ). A second peak is located around  $g = 2$ . Its intensity ( $I_g = 2$ ) decreases when the kaolinite cristallinity diminished. The intensity ( $I_g = 2$ ) is also influenced by small amounts of oxihydroxides.

Iron oxides or hydroxides are often associated with Fe<sup>3+</sup>-kaolinite causing an enlargement of the (Ig = 2) peak. A purifying treatment removing the free oxihydroxydes (Endrey, 1963) generally shows the disappearance of this (Ig = 2) signal. Then, the iron content of the material can be considered as structural.

*C. Range of aluminum and iron respectively in goethites, hematites, and kaolinites.*—In natural and synthetic goethites substitutions of Al<sup>3+</sup> for Fe<sup>3+</sup> range from 0 to 33 percent mole fraction of AlO(OH) (Benešlavsky, 1957; Simon, Lang, and Seidel, 1959; Thiel, 1963; Petit, Backer, and Herzog, 1964; Wefers, 1967; Janot and Gibert, 1970; Solymar, 1970; Janot and others, 1971; Bardossy, 1968, 1982).

In natural and synthetic hematites, the substitutions range between 0 and 15 percent mole fraction of Al<sub>2</sub>O<sub>3</sub> (Pouillard, 1950; Caillère, Gatineau, and Hénin, 1960; Shirane, Cox, and Roby, 1962; Bardossy, 1968, 1982).

In natural kaolinite, the mole fraction of Fe<sup>3+</sup>-kaolinite does not exceed 3 percent according to Janot, Gibert, and Tobias (1973), Rengasamy, Krishna-Murti, and Sarna (1975), Herbillon and others (1976), Fayolle (1979), Mendelovici, Yariv, and Villalba (1979), Didier (1983), and Cantinolle and others (1983). Furthermore, Herbillon and others (1976) and Mestdagh, Vielvoye, and Herbillon (1980) have shown that an increasing substitution of iron in kaolinite induces a diminution of cristallinity and an augmentation of disordering.

In conclusion, goethites and hematites can incorporate quite large amounts of aluminum in their structures. The mole fraction of AlO(OH) in goethites appears generally greater than the mole fraction of Al<sub>2</sub>O<sub>3</sub> admitted into hematites. Furthermore, the iron content in the kaolinite structure is significant but small. The degree of substitution of Al<sup>3+</sup> for Fe<sup>3+</sup> in goethites or hematites as well as the degree of substitution of Fe<sup>3+</sup> for Al<sup>3+</sup> in kaolinites is highly variable under natural conditions. Substitutions have undoubtedly a determinant role, even if it is not yet quantitatively appreciated, on the cristallinity and solubility of these minerals. Therefore, their stability fields have to be reconsidered in terms of solid solution equilibria.

#### DISTRIBUTION OF AL-HEMATITE AND AL-GOETHITE IN LATERITES

An abundant literature is devoted to the distribution of hematite and goethite in soils and weathering products such as bauxites, iron ferricretes, tropical kaolinitic alterations, and, more generally, laterites.

*A. Bauxites.*—In aluminum rich horizons and bauxites, gibbsite is generally the dominant aluminous mineral. As the formations get older boehmite progressively replaces gibbsite, and if burial or metamorphism takes place, even corundum may appear (Goldich and Berquist, 1948; Nia, 1968; Bardossy, 1982).

In most of the bauxites, Bardossy (1982) shows that hematite and goethite are often present and also that hematite dominates when boehmite is present. Furthermore, as described by Combes (1969) the classical bauxite parageneses are the following:

1. kaolinite and goethite are often closely related;
2. when gibbsite replaces kaolinite, goethite tends to be stable;
3. when boehmite, to a lesser extent, or diasporite, to a greater extent, appears, kaolinite, gibbsite, and goethite tend to disappear. Then, hematite dominates in the iron-rich phases.

Boulangé (1983) in a recent and extensive study of bauxites in the Ivory Coast, has clearly demonstrated that:

1. hematite and boehmite dominate in nodules and interglauclular plasmas of the initial aluminous ferricretes;
2. the amount of boehmite increases from the bottom to the top of the bauxitic profiles and becomes dominant in the pisolitic facies;
3. in the concretionary facies, boehmite tends to be more concentrated in the core of pisolites than in the surrounding plasma.

Cantinolle and others (1983) have analyzed the mineral assemblage of the Canonnettes bauxites (southeast of France) and have found that in the mottled kaolinitic bauxite, the crystallinity of kaolinite decreases, while iron content in kaolinite and aluminum content in the coexisting goethites increases together from the bottom to the top of the profiles.

In bauxitic formations, Norrish and Taylor (1961), Janot and Gibert (1970), Solymar (1970), Janot and others (1971), Mendelovici, Yariv, and Villalba (1979), and Bardossy (1982) have found substitutions of  $\text{Al}_2\text{O}_3$  in hematite ranging from 4 to 14 mole percent. Furthermore, they found substitutions of  $\text{AlO}(\text{OH})$  in goethite solid solutions, ranging between 5 and 33 mole percent. The last number represents to our knowledge the highest value found in natural samples (table 1).

In bauxitic profiles, the dehydrated minerals, hematite and boehmite, tend to be associated in the core of the pisolites, whereas the hydrated minerals, goethite and gibbsite, are formed together either in the initial or in the latest stages of the development of the profiles. The Al-content of goethite or hematite increases toward the top of the weathering formations.

*B. Gibbsite or kaolinite acid soils.*—According to Taylor and Graley (1967) hematite dominates in the upper parts of the old profiles. Therefore, the ratio hematite/goethite increases when the temperature rises, the humidity of the climate decreases, or the aridity increases, when the soil pH increases (Kampf and Schwertmann, 1983).

Aluminum content in hematite of oxisols analyzed by Schwertmann, Fitzpatrick, and Le Roux (1977) is reported in table 1. It is shown that the substitution is smaller in hematite than in goethite.

Fitzpatrick and Schwertmann (1982) have analyzed the aluminum content of goethites in strongly acid soils (gibbsitic soil or ferrallitic soils), moderately acid soils (kaolinitic soils or ferruginous tropical soils), and kaolinite bearing saprolites. Some geodes, mottles, or concretions were also considered. This extensive study shows clearly that the goethites of concretions, mottles, and kaolinitic, moderately acid soils are less substi-

tuted than those of kaolinitic saprolite and even less than those of strongly acid soils containing gibbsite (fig. 1).

*C. Ferricretes on lateritic profiles.*—In ferricretes, which are in a sense equivalent to bauxites, but, developed under seasonally contrasted tropical climate, hematite, goethite, kaolinite, and in many cases, quartz are often present but diversely distributed. The distribution of ferricretes as a function of time, climate, position in landscape, and nature of the parent material has been largely discussed over the past 50 yrs, among others by de Chetelat (1938), D'Hoore (1954), Maignien (1958), Millot (1964), Leprun (1971, 1972, 1979), Michel (1973) Nahon (1976), Nahon and Millot (1977), Boulet (1974), Nahon and others (1977), Didier (1983), Didier and others (1983), and Nahon (1985).

Typically, an old and well preserved lateritic weathering profile on granitic rock is represented in figure 2. Five different horizons are distinguished (Tardy, 1969; Nahon, 1976):

1. The fresh parent granitic rock.
2. The saprolite, in which, together with smectite and kaolinite, some feldspars or primary micas remain unaltered.

TABLE 1  
Average aluminum content in goethite of various origins

Formation	References	Mineral	Composition		Associated minerals
			AlO(OH)	FeO(OH)	
			Al <sub>2</sub> O <sub>3</sub>	Fe <sub>2</sub> O <sub>3</sub>	
Bauxite	Morrish and Taylor (1961)	Goethite	14.5	85.5	Kaolinite, gibbsite
Bauxite	Bardossy (1968)	Goethite	17.0	83.0	Kaolinite, gibbsite, boehmite
Bauxite	Bardossy (1968)	Hematite	3.0	97.0	Kaolinite, gibbsite, boehmite
Bauxite	Janot and Gilbert (1970)	Hematite	12.0	88.0	Boehmite
Bauxite	Janot and Gilbert (1970)	Hematite	5.0	95.0	Kaolinite, gibbsite (or boehmite)
Bauxite	Janot and Gilbert (1970)	Hematite	5.0	95.0	Kaolinite, gibbsite (or boehmite)
Bauxite	Solymar (1970)	Goethite	22.5	77.5	No records
Bauxite	Janot and others (1971)	Goethite	10.0	90.0	Kaolinite, gibbsite
Bauxite	Janot and others (1971)	Goethite	15.0	85.0	Kaolinite, gibbsite
Bauxite	Janot and others (1971)	Goethite	12.0	88.0	Kaolinite, gibbsite
Bauxite	Janot and others (1971)	Hematite	12.0	88.0	Diaspore, corundum
Bauxite	Janot and others (1971)	Goethite	25.0	75.0	Kaolinite, gibbsite, boehmite
Bauxite	Janot and others (1971)	Hematite	8.0	92.0	Kaolinite, gibbsite, boehmite
Bauxite	Fitzpatrick and Schwertmann (1982)	Goethite	24.0	76.0	Kaolinite, gibbsite
Laterite	Muller and others (1981)	Goethite	15.0	85.0	Kaolinite, gibbsite
Ferricrete	Nahon (1976)	Goethite	20.0	80.0	Kaolinite, quartz
Ferricrete	Nahon (1976)	Hematite	13.0	87.0	Kaolinite, quartz
Ferricrete	Nahon and others (1979)	Goethite	15.0	85.0	Kaolinite
Ferricrete	Nahon and others (1979)	Hematite	15.0	85.0	Kaolinite
Ferricrete	Leprun (1979)	Goethite	20.0	80.0	Kaolinite, quartz
Ferricrete	Leprun (1979)	Goethite	17.0	83.0	Kaolinite, quartz
Ferricrete	Leprun (1979)	Goethite	17.0	83.0	Kaolinite, quartz
Ferricrete	Fitzpatrick and Schwertmann (1982)	Goethite	9.0	91.0	No records
Strongly acid soils	Fitzpatrick and Schwertmann (1982)	Goethite	23.0	77.0	No records
Moderately acid soils	Fitzpatrick and Schwertmann (1982)	Goethite	7.0	93.0	No records
Saprolites	Fitzpatrick and Schwertmann (1982)	Goethite	15.0	84.0	No records
Mottles	Fitzpatrick and Schwertmann (1982)	Goethite	6.0	94.0	No records
Oxisols	Schwertmann, Fitzpatrick and Le Roux (1977)	Hematite	8.2	91.8	No records
Oxisols	Schwertmann, Fitzpatrick and Le Roux (1977)	Hematite	5.5	94.5	No records
Red soils	Torrent, Gunzman and Parra (1982)	Goethite	10.0	90.0	Kaolinite
Red soils	Torrent, Gunzman and Parra (1982)	Hematite	3.0	97.0	Kaolinite




Fig. 1. Aluminum content in goethites from various origins (from Fitzpatrick and Schwertman, 1982).

3. The lithomarge, in which all primary minerals: that is plagioclases, K-spars, biotites, and amphiboles are entirely transformed into kaolinite. The initial structure of the rocks is preserved, and the isovolumetric chemical balance shows a strict conservation of aluminum and iron. Furthermore, iron is localized as goethite and located at the places previously occupied by ferro-magnesian primary minerals. Beside these pink or red-dish-colored places the kaolin-transformed feldspars remain white, so that, at this level, the mobilization of iron is not yet observed. The differences

in color are determined, prior to the alteration, by the initial distribution of iron in primary minerals. The iron and color distribution are referred to as litho-relictual.

The lithomarge is then a formation in which quartz, kaolinite, and goethite coexist, all preserved along an important thickness, as long as its situation remains below the water table level.

In the saturated zone permanently located under the water table, hematite is absent, and no concretion formation is observed (Nahon, 1976; Muller and others, 1981).

4. The mottled formation overlies the lithomarge above the water table level, in the unsaturated zone. Here, progressively the initial structure of the parent rock is effaced or obliterated. Two major pedoturbations affect the mottled formation. First of all, this horizon is invaded by




Fig. 2. Schematic representation of a common lateritic profile and its overlying ferricrete.

a network of channels and tubular voids of large diameter (cm) in which both detrital and neoformed kaolinite can accumulate. Furthermore, iron appears now as mobilized, that is, leached from the areas located around large pores (mm) and concentrated in clay-rich areas, forming mottles, precursors of concretions. The secondary kaolinite accumulated in large tubular voids is also progressively enriched in iron so that iron concentrates in areas consisting of fine grained material characterized by a small pore size ( $\ll 0.1$  mm). The original lithostructure is now progressively replaced toward the top of the profiles by a newly formed pedostructure. Here, in the mottles, goethite dominates, but hematite appears in small percentages. When mottles evolve into concretions, hematite replaces goethite.

5. The mottled formation is covered by the ferricrete *sensu stricto*. This horizon, generally a few meters thick, consists of an iron accumulation which can be more than 20 percent  $\text{Fe}_2\text{O}_3$ . In the middle part, cemented nodules are abundant. To the maximum in concretion formation correspond the maximum hardness and the maximum hematite development. It corresponds also to the complete replacement of the kaolinitic host by an aluminum rich hematite. Toward the top, rings of goethite appear around hematitic nodules as a secondary degradation process taking place at the edges of the large pores. The secondary goethites are formed by rehydration of Al-hematites. They are highly aluminous and become poorer and poorer in aluminum as long as the degradation process takes place. The end product is a pure goethite. Close to the top, the cohesion diminishes, and nodules have a tendency to separate from each other so that iron ferricretes are very often covered by gravels made of individual and deconsolidated nodules. Thus, the superficial gritty horizon is a result of ferricrete dismantling (Nahon, 1976). This is schematized in figure 2.

Dissolved iron moves downward from the top and accumulates in the bottom part of the crust. This movement provides iron for the concretion and nodule formation. Progressively aluminum and silicon are removed from the ferricrete; kaolinite dissolves, quartz grains are highly corroded, whereas hematite and goethite are retained and become the dominant minerals.

In conclusion, when, at the bottom of the ferricretes, goethites and hematite form together, goethite is generally of a low Al-content. This is interpreted as evidence that goethite precipitates in the large pores at the contact of the quartz grains, for instance, and that hematite forms in the smallest pores of kaolinite material which is progressively replaced and which yields the amounts of aluminum required for the Al-hematite formation.

Therefore, the highest aluminum contents in hematites are found together with the maximum development of the nodules and the maximum dissolution of kaolinite. Furthermore, the highest aluminum contents in goethites are found accompanying the first stages of rehydration of Al-hematites together with the dismantling of the nodular horizon.



*D. Conclusion: field observations and distribution of Al-goethite and Al-hematite.*—As seen above, hematite and goethite are often associated in tropical red soils: laterites, ferricretes, and bauxites. Hematite tends to form on the top of the profiles and in the small size pores of the unsaturated zone. Goethite tends to form at the bottom of the profiles under the water table zone. It develops also in edges of the large pores and channels which are occasionally visited by sudden water discharges. Obviously, the hematite-goethite distribution is regulated by water activity.

In addition to these general observations, it should be remarked that goethite does not replace minerals, and especially kaolinite, with which it is associated. On the contrary, hematite, when it forms instead of goethite, replaces kaolinite which dissolves. This is, in our opinion, the reason hematite concretions and nodules can be formed when the water activity of the system diminishes (fig. 3).

From natural observations, it can also be concluded that the aluminum contents of goethite and hematite should be regarded as a function of the nature of other minerals with which they are associated.

For the 183 samples analyzed by Fitzpatrick and Schwertmann (1982) and other data presented in table I, one may underline two major facts:

1. In goethites of red soils, the substitution rates diminish when the climate becomes less humid or more arid and percolating solutions are richer in silica (Tardy, 1971).

2. Compositions range from 2 to 20 mole percent of AlO(OH) in goethites of ferricretes and other soils in which kaolinite is present. Compositions range from 18 to 27 mole percent in goethites associated with gibbsite of bauxites.




Fig. 3. Formation and evolution of a mottle into a nodule and into a pseudo-pisolite (after Nahon, 1976).

Thus, the substitution rate of  $\text{AlO}(\text{OH})$  into  $\text{FeO}(\text{OH})$  is lower in ferricretes than in bauxites, as generally admitted by Nahon (1976), Nahon and others (1979), Leprun (1979) in laterites of Senegal and Ivory Coast. Hematite is generally substituted less than goethite, and both appear less aluminous when kaolinite and quartz are present than when gibbsite forms.

Obviously, the aluminum contents of goethites and hematites are controlled by the activity of aluminum in solution which is in turn determined by the solubility of aluminous minerals, that is, gibbsite and kaolinite.

It turns out that the relative stability fields of goethite and hematite may depend on their degree of Al-substitution which are dependent on the stability of the other aluminous minerals with which they are associated. The conditions of formation of goethite and hematite may be ultimately determined by the activity of silica in solution and by the activity of water in pores in which these minerals form.

#### FACTORS CONTROLLING THE FORMATION OF GOETHITE AND HEMATITE

A large number of works have been devoted, over the last 20 yrs, to the stability of iron oxides and hydroxides in the laboratory and under natural conditions (Schwertmann, 1965; Schmalz, 1959, 1968; Berner, 1969; Bischoff, 1969).

Three kinds of factors are generally considered: grain size effects, kinetics of precipitation, and equilibrium conditions involving the activity of water. Because of its importance, water activity is discussed in a separate section. In the present section only grain size effects and kinetics are treated.

*A. Grain size effect.*—Langmuir (1971) has treated this problem extensively and has shown that the relative stability of Al-free goethite and hematite in water is strongly dependent on the particle size of the minerals involved. Because the heats of dissolution as a function of particle size are different for goethite and hematite, the equilibrium reaction energy is also a function of the dimensions of the crystals. When crystals of hematite and goethite are of the same size, hematite appears to be more stable than goethite. This is also, of course, true when hematite crystallizes in a larger size. However, the opposite is observed when hematite forms in a smaller grain size than goethite.

In the cases of ferricretes, two situations are observed. In the first steps of the formation of mottles and concretions, hematite and goethite are both of a small size, and there is difficulty in saying which is coarser than the other. During the last steps of nodule dismantling, only, the secondary goethite appears in quite well formed crystals of about  $1\ \mu\text{m}$  diameter.

Because hematite and goethite in concretions and nodules never show large crystals but only very tiny particles of about  $100\ \text{\AA}$  in diameter, it seems clear that conditions governing their stability must be related more to dehydration and water activity and less to particle size effects.

*B. The kinetic interpretation.*—The hematite and goethite formation mechanism in soils has been discussed by Schwertmann and Taylor (1977). Kampf and Schwertmann (1983) have written that “Iron from silicates and other primary sources released into solution may lead to a direct crystallization of goethite if the solubility product of goethite, but not the much higher one of ferrihydrite, is exceeded. If the solubility product of ferrihydrite is exceeded however, ferrihydrite will be formed and may then transform either to hematite via a dehydration, or alternatively to goethite via dissolution of ferrihydrite”. One may understand here that the primary factor that governs the rate of dissolution versus precipitation of iron is kinetic. Then, these authors continue their interpretation and state that factors favoring hematite over goethite formation are those that favor ferrihydrite formation such as a rapid release of Fe and a low concentration of organic compounds which complex Fe, thereby allowing a somewhat higher concentration of inorganic Fe<sup>3+</sup> ions. Conversely organic compounds favor goethite formation, because the Fe-complex formation prevents the attainment of ferrihydrite saturation.

In addition, once ferrihydrite is formed, hematite is favored over goethite with increasing temperature because dehydration is involved in the ferrihydrite → hematite transformation.

This interpretation is based fundamentally on laboratory experiments and statistical correlations between mineralogical data and a set of environmental factors. Kämpf and Schwertmann (1983) themselves have recognized: “the hematite-goethite relationship cannot be elucidated from our study, however, because of the strong intercorrelation among factors considered such as temperature, excess of moisture and even soil organic matter and pH”.

Berner (1969), reviewing the stability conditions of hematite in water, has pointed out that, although hematite is more stable than goethite, kinetic phenomena may be the reason for the non-spontaneous formation of hematite in water. Slow transformation of original metastable goethite should take place during diagenesis, and this is the probable reason why goethite is absent in buried sediments ranging from Tertiary to Paleozoic in age (Fischer, 1963). Langmuir (1969) is in agreement with Berner's (1969) view that fine-grained goethite has no thermodynamic stability relative to well crystallized hematite. Then, for kinetic reasons, goethite is more common than hematite, and also for kinetic reasons, once hematite has appeared, it does not rehydrate to form coarse goethite.

However, natural observations made on ferricrete suggest that besides the kinetic interpretation, the distribution of goethite and hematite can also be understood in terms of thermodynamic equilibria in which the chemical potential of water is involved.

#### ACTIVITY OF WATER, PORE SIZE EFFECT, AND CONCRETION FORMATION

*A. Activity of water ( $a_w$ ).*—By definition, the activity  $a_i$  of any substance, subscribed as  $i$  (Garrels and Christ, 1965), is defined as:

$$a_i = f_i/f_i^\circ$$

in which  $f_i$  and  $f_i^\circ$  stand respectively for the fugacity of the vapor above the substance, and  $f_i^\circ$  the fugacity of the vapor when the substance is in its standard state. For water vapor in the atmosphere, at room temperature, and under normal atmospheric total pressures, fugacities can be regarded as equal to partial pressures, so that:

$$a_w = p/p_o = \text{R.H.}/100$$

in which  $p$  and  $p_o$  stand respectively for the water partial pressure above the soil and the water partial pressure of the corresponding water-saturated air. R.H. (percent) is then the relative humidity of the air measured at a given temperature and pressure.

In porous media and in the unsaturated zone of soils, the activity of capillary water, retained behind the concave meniscii, is equal, if equilibrium is reached, to the relative humidity of the air imposed by climatic conditions (see Hillel, 1971; or Sposito, 1981).

When rainfall ceases, the water percolation stops, and after drainage and evaporation, water is retained in pores of progressively smaller and smaller radii and consequently behind smaller and smaller air-water interface meniscii. As a result, water activity also diminishes progressively (see Bourrié and Pédro, 1979; Dandurand and others, 1982). The Laplace's formula allows calculation of the pressure difference between the two sides of the air-water interface as a function of  $\sigma$ , the interfacial tension (72.7 dyn/cm at 25°C for pure water) and  $r$  the radius of the circular section of the pore:

$$\Delta P = -2 \sigma / r$$

The change in the chemical potential of the water is a function of its molar volume ( $\bar{V}$ ) and the pressure difference ( $\Delta P$ ) between each side of the meniscus:

$$(\mu - \mu_o) = RT \ln a_w = \bar{V} \cdot \Delta P = -\bar{V} \cdot 2 \sigma / r$$

In all cases of capillary waters, these terms are all negative because  $\Delta P$  is negative. The scale of correspondence between the size of cylindrical shaped pores and the activity of water is given in table 2. It is interesting to remark that a relative humidity of 0.5, commonly observed during the dry season of tropical climates, corresponds to a pore diameter of about 30 Å which is precisely the diameter of pores provided by the edges of kaolinite grains.

*B. Hydration-dehydration isotherms and the pore diagram of kaolinite.*—Petrographic observations have shown that hematite concretions form preferentially in sites previously enriched in kaolinite and not in sites previously leached and made of quartz skeleton only.

The question that emerges concerns the pore size distribution offered by kaolinite aggregates compared to that shown by quartz grain assemblages. The pore size distribution can be calculated from adsorption or desorption isotherms, assuming that the pore shape is known. In the literature, most of the BET surface areas and pore size distributions are

TABLE 2

Correspondence between the diameter of the meniscus (or capillary) and the equilibrium water activity

Pore diameter		Activity of water
d	= 30 $\mu\text{m}$ ( $3 \times 10^{-5}\text{m}$ )	$a_w = 0.99993$
d	= 3 $\mu\text{m}$ ( $3 \times 10^{-6}\text{m}$ )	$a_w = 0.9993$
d	= 0.03 $\mu\text{m}$ ( $3 \times 10^{-8}\text{m}$ )	$a_w = 0.9312$
d	= 0.003 $\mu\text{m}$ ( $3 \times 10^{-9}\text{m}$ )	$a_w = 0.4903$
d	= 0.0003 $\mu\text{m}$ ( $3 \times 10^{-10}\text{m}$ )	$a_w = 0.0009$
$d(\text{m}) = -0.0921 \cdot 10^{-8} / \log a_w$		

based on nitrogen adsorption, and only a few are derived from water adsorption or desorption. Depending on the type of molecule used (N<sub>2</sub> or H<sub>2</sub>O) the results may be different particularly concerning pores of very small size (several Å). Because natural processes occur in water media, we have performed and selected a series of measurements and calculations devoted to water adsorption or desorption on kaolinite.

Kaolinite isotherms have been presented by Keenan, Mooney, and Wood (1951), Torrence-Martin (1958), and Kehres (1983). An example taken from Kehres (1983) is shown in figure 4. It concerns a Georgia kaolinite. A dehydration isotherm is a curve obtained by plotting  $\theta$ , the water content (grams of H<sub>2</sub>O per gram of dry clay) as a function of water activity ( $a_w$ ). The activity of water is defined as the ratio  $a_w = p/p_0$ ;  $p$  is the equilibrium partial pressure of vapor at a given stage of hydration, and  $p_0$  is the equilibrium partial pressure at saturation (pure liquid water in standard state).

From the data of the dehydration isotherm (right side of fig. 4), two kinds of results are derived. The first concerns the BET surface area, and the second the pore size distribution. The BET surface area (Brunauer




Fig. 4. Dehydration isotherm and BET surface area of kaolinite (after Kehres, 1983).

and Emmett, 1935, 1937; Brunauer, Emmett, and Teller, 1938) is obtained by plotting  $p/(\theta(p_0 - p))$  as function of  $a_w = p/p_0$  (left side of fig. 4). Between  $p/p_0 = 0.1$  and  $0.4$  the function is linear, so that:

$$p/(\theta(p_0 - p)) = \alpha (p/p_0) + \beta$$

and the BET theory says that:

$$\alpha = \frac{C-1}{V_m C} \text{ and } \beta = \frac{1}{V_m C}$$

By knowing  $\alpha$  and  $\beta$ , it is possible to calculate  $C$ , a function of the energy of adsorption, and  $V_m$ , the volume of a monolayer adsorbed. The surface area  $A$  is easily calculated from the monolayer volume  $V_m$ . The Georgia kaolinite studied presents a BET surface area of about  $35 \text{ m}^2/\text{g}$ . Clay minerals such as disordered kaolinites may present a quite large surface area mostly due to defects of layer stacking. Then, an important part of the surface area is created along the edges, in pores of small size.

The pore size distribution can be calculated by following two methods. The first is based on the application of Kelvin's formula for cylindrical pores (Pierce, 1953). Another method is used here in order to take into account that clay minerals are planar layers and that pores should have planar walls instead of cylindrical ones (Delon and Dellyes, 1967).

Pierce's method for pores of cylindrical shape is based on the following approach. For a given  $p/p_0$ ,  $\theta$  is the volume of water per gram of dry clay trapped in pores of a diameter smaller than  $d$ .

To an increment  $\Delta(p/p_0)$  corresponds a volume change  $\Delta\theta$  and an internal area, change  $\Delta A$ . The average pore diameter  $\bar{d}$  concerned is included between  $d$  and  $d + \Delta d$ , and one defines a distribution function  $S(\bar{d})$ :

$$S(\bar{d}) = \Delta A / \bar{d}, \text{ for internal } \Delta(p/p_0);$$

$S(\bar{d})$  is in fact constant up to a given length  $l$  and in the case of a cylinder;

$$S(\bar{d}) = \pi \bar{d} l / d = \pi l.$$

The Delon-Dellyes's (1967) method considers pores of parallel planar faces. The Kelvin's relationship has a different formulation, but the pore distribution function,  $S(\bar{d})$  is calculated similarly:

$$S(\bar{d}) = \Delta A / \bar{d}$$

$\bar{d}$  is the average distance between the planar face concerned by the corresponding surface area  $\Delta A$ . The numbers given in ordinate axes in figure 5 correspond in fact to additions of pore length  $10^{20} \text{ \AA}$  per gram of clay calculated per intervals of radius or diameter ( $\Delta d$ ), derived from intervals of water activity  $\Delta(p/p_0)$ .

The distribution function  $S(\bar{d})$  calculated from Pierce's method corresponds to a summation of real pore length, because these pores are considered as cylindrical. The distribution function  $S(\bar{d})$  calculated from

Delon-Dellyes's method is derived from the same equation and also has the dimension of a length [L]. However, it does not correspond to a real summation of pore length, because the pores are supposed, in this case, to be of a planar shape.

The two diagrams of figure 5, characterizing a Georgia kaolinite studied by Kehres (1983), present almost the same distribution function. The maximum of the length distribution is obtained for a pore radius of about 20 Å (let us say 21 Å) or a pore diameter of about 40 Å (let us say 42 Å) which corresponds to an average of six 7 Å-kaolinite layers.

By comparison between kaolinite aggregates and quartz powder of about the same grain size (2.4 µm) the surface area of a disordered kaolinite is, in general, much greater (35 m<sup>2</sup>/g instead of 5 m<sup>2</sup>/g), and the pores of small size (< 50 Å in diam) are much more numerous in kaolinite than in quartz.

*C. Porosity differences between kaolinite cutans and quartz sands.—*

In tropical soils, there are domains enriched in kaolinite coexisting with zones in which quartz sand dominates. The former correspond to areas of parent material initially enriched in biotite and amphibole or to cutans made by illuviation and accumulation of fine grained material. The latter correspond to areas of parent material initially rich in quartz or to eluviated zones from which kaolinite has been leached.

At low water activities, kaolinitic cutans may retain much more water than quartz sand, simply because the proportion of small pores is quite high at the edges of kaolinite crystals and extremely low at the contact between quartz grains. At high water activities, the contrary is observed,




Fig. 5. Pore diagram of kaolinite (after Kehres, 1983).

and the porosity of quartz sand filled by water could be very large. It seems not unreasonable to expect that solutions that become saturated with respect to iron minerals should precipitate goethite in water of high activity at the contact with large sized pores and hematite in water of low activity in very small sized pores. Therefore, another mechanism should take place and explain formation of goethitic mottles and the development of the hematitic nodules and concretions.

*D. Mottles, concretions, and nodule formation.*—One of the most remarkable facts in the process of a ferricrete formation is that iron is leached from large sized pores and accumulates in small sized pores, that is, moves from sand to clay.

This process starts to take place above the water level and develops farther toward the soil surface. Under the water table, in the permanently wet horizons, no migrations are observed.

Above the water table level, mottles begin to form, and the first mineral that concentrates is goethite. Farther upward, hematite follows goethite, and nodules develop from mottles. In both cases, iron moves from the outer part to the inner part of the concretion. The driving force of this migration and accumulation seems to be the initial difference in the size of the pores which tends to be accentuated as the concretion develops. The interpretation of this phenomenon was proposed by Tardy and Monnin (1983) who have calculated the mineral solubility differences as functions of the size of the pores in which they precipitate. Briefly, these authors have proposed that an aqueous solution at saturation with respect to a given mineral in a pore of a large size can be oversaturated with respect to the same mineral in a pore of small size. In other words, water circulating in large channels dissolves minerals, becomes saturated with respect to them, and fills the pores of small size. After drainage, the small sized pores remain filled by water retained behind concave menisci of small radii. In these conditions, the solutions become oversaturated with respect to the corresponding minerals which precipitate and tend to fill the porosity which again decreases in size; thus, the initial process is perpetuated.

Two reasons may be responsible for these changes in the saturation state of solutions. The first corresponds to the changes in the activity of solids as a function of the shape of their surface. An activity of unity characterizes pure phases presenting infinite and flat surfaces. An activity greater than unity characterizes generally pure phases presenting convex interfaces behind which the pressure is locally greater than in the external domain. In such cases, solids are more soluble than they are in their standard state. An activity smaller than unity characterizes generally pure phases presenting concave interfaces behind which the pressure is locally smaller than in the external domain. Solids are, then, less soluble in such cases than they are in their standard state (Wollast, 1971). This is the reason a solid surface presenting needles and cracks evolves naturally and becomes flat: needles dissolve, and cracks are filled. Iler (1979) has shown that such a mechanism is responsible for the evolution of silica-gels, where-



as Tardy and Monnin (1983) and Dandurand and others (1982) have proposed that it is responsible for the formation of silica concretions and for silcrete formation.

The second phenomenon acting in the same way takes place when, after drainage, water remains, behind concave menisci, in pores of small size. Here, in water, the pressure is smaller than in the atmosphere, and the activity of water is smaller than unity. Tardy and Monnin (1983) have proposed that minerals that are more soluble when pressure increases would be less soluble when pressure decreases, that is, behind concave menisci. In fact, Dandurand and others (1982) have verified that aqueous silica concentration at equilibrium with respect to amorphous silica diminishes when the pores concerned become smaller and smaller.

Iron minerals, such as goethite and hematite, also follow these rules and are both more soluble when particle size decreases and when pressure increases.

These different mechanisms all act in favor of a concentration of goethite and hematite in pores of small size and in favor of the mottle formation concretioning and nodulation. When the size of pores in which the precipitation reaction takes place, that is, when the activity of water is small enough, goethite no longer forms, and hematite becomes the dominant mineral. A decrease of the activity of water is not only responsible for concretion formation but also for the nature of the mineral formed.

#### EQUILIBRIUM REACTIONS AS FUNCTIONS OF WATER AND SILICA ACTIVITIES


Because goethite and hematite are respectively hydrated and dehydrated minerals, their equilibrium boundary is first of all dependent on the activity of water. Also, because Al-goethite is a solid solution of diaspora and goethite and Al-hematite a solid solution of corundum and hematite, the chemical compositions of the coexisting solid solutions are controlled by water activity in the system  $\text{Al}_2\text{O}_3\text{-Fe}_2\text{O}_3\text{-SiO}_2\text{-H}_2\text{O}$ . Furthermore, because kaolinite interferes in the control of Al ions in solution, the chemical compositions of goethite and hematite are also dependent on the activity of dissolved silica.

*A. Solubility products of minerals.*—In the following sections we have used conventional aqueous species for writing equilibrium equations and calculating solubility products of the following minerals:




and

$$\log [\text{Al}^{3+}]/[\text{H}^{3+}] + 2 \log [\text{H}_2\text{O}] = \log K_{\text{sp}} \text{ diaspore}$$




and

$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 + 1.5 \log [\text{H}_2\text{O}] = \log K_{\text{sp}} \frac{1}{2} \text{ corundum}$$



and

$$\log K_{\text{sp}} \text{ gibbsite} = \log [\text{Al}^{3+}]/[\text{H}^+]^3 + 3 \log [\text{H}_2\text{O}]$$



$$\text{and} \quad (3)$$

$$\log K_{sp} \text{ 1/2 kaolinite} = \log [\text{Al}^{3+}]/[\text{H}^+]^3 + \log [\text{SiO}_2] + 2.5 \log [\text{H}_2\text{O}]$$

$$\text{goethite: } \text{FeO}(\text{OH}) + 3\text{H}^+ \rightleftharpoons \text{Fe}^{3+} + 2 \text{H}_2\text{O} \quad (4)$$

and

$$\log [\text{Fe}^{3+}]/[\text{H}^+]^3 + 2 \log [\text{H}_2\text{O}] = \log K_{sp} \text{ goethite}$$

$$\text{1/2 hematite: } \text{1/2 Fe}_2\text{O}_3 + 3\text{H}^+ \rightleftharpoons \text{Fe}^{3+} + 1.5 \text{H}_2\text{O} \quad (5)$$

and

$$\log [\text{Fe}^{3+}]/[\text{H}^+]^3 + 1.5 \log [\text{H}_2\text{O}] = \log K_{sp} \text{ 1/2 hematite}$$

$$\text{quartz: } \text{SiO}_2 \rightleftharpoons \text{SiO}_{2(\text{aq})} \quad (6)$$

and

$$\log [\text{SiO}_2] = \log K_{sp} \text{ quartz} \quad (7)$$

*B. A thermodynamic and solubility product data set.*—The Gibbs free energies of formation and the solubility products of minerals used are reported in tables 3 and 4. They include data presented by Robie and Waldbaum (1968), Langmuir (1969), Naumov, Ryzhenko, and Khodakovsky (1971), Berner (1971), Hem (1970), Hem and Roberson (1967), Hem and Lind (1974), Hem and others (1973), Parks (1972), Ulbrich and Merino (1974), Helgeson and others (1978), Hemingway, Robie, and Kittrick (1978), Robie, Hemingway, and Fischer (1978), Khodakovsky (1983, personal commun.).

The solubility product values of minerals and of the endmembers involved in the solid solutions were discussed by Fritz and Tardy (1974), Fritz (1981), Tardy and Fritz (1981), Tardy (1982), and Didier (1983). In fact, when looking at the stability field of minerals in nature, it appears that the solubility product values listed in table 4 are not all internally consistent and are not capable of illustrating correctly the observed mineral associations and the Al/Fe ratios in goethite and hematite natural solid solutions.

We started from the solubility products of gibbsite (8.205) and kaolinite (7.41) chosen by Fritz and Tardy (1974). These values are compatible with the fact that quartz and kaolinite coexist and have been chosen over the years as the “best values” for kaolinite and gibbsite formed in laterites.


The solubility product of boehmite is derived from the data of Robie, Hemingway, and Fischer (1978), by subtracting from our gibbsite  $K_{sp}$  value the same difference as found in column 8 of table 4:

$$\log K_{sp} \text{ boehmite} = \log K_{sp} \text{ gibbsite} + 0.14 = 8.205 + 0.14 = 8.345 \quad (8)$$

Other values were adapted in order to be compatible with natural observations and the ideal solid solution model. These numbers are reported in the last columns of tables 3 and 4. It seems clear that goethite, hematite, and diasporite solubility products are higher than those currently

quoted in the literature. This is supported by the fact that the solid solutions found in nature are crystallizing in a very small size and should be more soluble than the corresponding well crystallized minerals.


*C. Gibbsite-kaolinite equilibrium.*—The solubility products of kaolinite and gibbsite are written as follows:



with

$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 = + 3.705 - \log [\text{SiO}_{2(\text{aq})}] - 2.5 \log [\text{H}_2\text{O}] \quad (9)$$

and



with

$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 = 8.205 - 3 \log [\text{H}_2\text{O}] \quad (10)$$

in which the terms between brackets designate activities of substances. At equilibrium between the two minerals, one has:

$$\log [\text{SiO}_{2(\text{aq})}] = - 4.5 + \frac{1}{2} \log [\text{H}_2\text{O}] \quad (11)$$

When water activity is equal to one, it becomes:


$$\log [\text{SiO}_{2(\text{aq})}] = - 4.5 \quad (12)$$

If the water activity diminishes, the activity of silica defined at equilibrium also diminishes and kaolinite becomes more stable. This has been already pointed out by Sposito (1981), Pédro and Delmas (1980), and Tardy (1982) and explains why at the same level of a profile kaolinite remains stable in the finely porous materials while gibbsite is stabilized at the edges of channels and large cavities.

*D. Al-goethite-kaolinite and Al-hematite-kaolinite equilibria.*—When kaolinite coexists with Al-goethite or Al-hematite, the equilibrium reactions should involve diaspore and corundum components. An Al-goethite ( $\text{Al}_x \text{Fe}_{1-x}\text{O}(\text{OH})$ ) is in fact a solid solution of  $x \text{AlO}(\text{OH})$  and  $(1-x)\text{FeO}(\text{OH})$ , corresponding to  $x$  diaspore and  $(1-x)$  goethite mole fractions.

An Al-hematite ( $\frac{1}{2} (\text{Al}_y \text{Fe}_{1-y})_2\text{O}_3$ ) is in fact a solid solution of  $y(0.5 \text{Al}_2\text{O}_3)$  and  $(1-y)(0.5 \text{Fe}_2\text{O}_3)$ , corresponding to  $y$  0.5 corundum and  $(1-y)$  0.5 hematite. Equilibria among solid solutions can be written by using partial equilibrium conditions (Fritz, 1981).

The solubility product of diaspore is written as follows:



$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 = 8.949 - 2 \log [\text{H}_2\text{O}] \quad (13)$$

When  $\text{AlO}(\text{OH})$  is involved in Al-goethite solid solution ( $(\text{Al}_x \text{Fe}_{1-x})\text{O}(\text{OH})$ ),  $x$ , being the diaspore mole fraction and  $\lambda_x$  the corresponding activity coefficient, one has:

$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 = 8.949 + \log x + \log \lambda_x - 2 \log [\text{H}_2\text{O}] \quad (14)$$

If the solid solution is ideal,  $\lambda_x = 1$ .

TABLE 3

Gibbs free energies of formation of minerals and aqueous species used. Selected data in this work are listed in column (11)

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
$\text{Fe}^{3+}$	- 10.54	- 17.87	- 10.54	---	---	---	- 4.60	- 4.60	- 4.60	---	- 4.60
$\text{Fe}_2\text{O}_3$	- 743.62	- 742.41	- 740.98	---	---	---	- 745.40	- 742.68	- 728.07	---	- 711.113
$\text{FeO}(\text{OH})$	---	- 490.36	- 488.27	---	---	---	---	- 488.55	- 483.471	---	- 474.305
$\text{Al}^{3+}$	- 485.34	- 492.00	- 481.16	- 481.16	- 485.34	- 485.34	- 489.36	- 489.40	- 489.90	- 489.40	- 489.400
$\text{Al}_2\text{O}_3$	- 1581.89	- 1557.28	- 1581.97	---	- 1582.39	---	- 1568.25	- 1582.23	- 1597.37	- 1581.08	- 1580.406
$\text{AlO}(\text{OH})$ (d)	---	- 921.32	- 914.62	---	- 918.39	- 916.50	- 913.78	- 922.00	- 921.30	- 920.80	- 912.702
$\text{AlO}(\text{OH})$ (b)	---	---	---	---	- 915.04	- 907.30	---	- 918.40	---	- 915.30	- 916.149
$\text{Al}(\text{OH})_3$	- 1144.28	- 1156.89	- 1144.32	- 1139.30	- 1151.85	- 1147.08	- 1155.48	- 1154.89	- 1153.98	- 1154.50	- 1154.138
$\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$	- 3777.61	- 3778.15	- 3776.90	- 3753.05	---	- 3779.95	- 3789.08	- 3799.36	- 3789.64	---	- 3789.374
$\text{SiO}_2$ (q)	- 856.25	- 856.51	- 856.05	- 856.88	---	- 856.21	- 856.25	- 856.29	- 856.29	---	- 856.29
$\text{SiO}_2$ (aq)	- 842.24	- 833.70	- 834.37	- 833.95	---	- 833.37	- 833.41	- 833.72	- 833.72	---	- 833.459
$\text{H}_2\text{O}$	- 237.19	- 237.19	- 237.19	- 237.19	- 237.19	- 237.19	- 237.19	- 237.14	- 237.19	---	- 237.190

(1) Robie, and Waldbaum (1968)

(2) Naumov, Ryzhenko, and Khodakovsky (1971)

(3) Berner (1971)

(4) Hem (1970); Hem, and Lind (1974); Hem, and Roberson (1973); Hem, and others (1973)

(5) Parks (1972)

(6) Ulbrich and Merino (1974)

(7) Helgeson (1978)

(8) Hemingway, Robie, and Kittrick (1978); Robie, Hemingway, and Fischer (1978); Hemingway, and Robie (1977a,b)

(9) Tardy (1982)

(10) Khodakovsky (1983)

(11) Proposed in this study

TABLE 4

Solubility products  $\log (K_{sp})$  derived from the Gibbs free energies of table 3. The values proposed in this study are given in column (11). See legend to table 3 for references corresponding to each column

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
Goethite $\text{FeO}(\text{OH})$	---	- 1.331	- 0.586	---	---	---	- 4.323	- 1.696	- 0.806	---	+ 0.819
Hematite $\frac{1}{2} (\text{Fe}_2\text{O}_3)$	- 0.962	- 0.425	- 0.729	---	---	---	- 2.161	- 1.943	- 0.663	---	+ 0.846
Diaspore $\text{AlO}(\text{OH})$	---	+ 7.890	+ 7.169	---	+ 7.239	+ 7.570	+ 8.922	+ 7.318	+ 7.324	+ 7.528	+ 8.949
Boehmite $\text{AlO}(\text{OH})$	---	---	---	---	+ 7.826	+ 9.180	---	+ 7.931	---	+ 8.474	+ 8.345
Gibbsite $\text{Al}(\text{OH})_3$	+ 9.218	+ 8.177	+ 8.480	+ 9.359	+ 7.892	+ 8.728	+ 7.961	+ 8.071	+ 8.230	+ 8.140	+ 8.205
Corundum $\frac{1}{2} (\text{Al}_2\text{O}_3)$	+ 11.721	+ 11.721	+ 8.519	---	+ 8.746	---	+ 10.688	+ 9.451	+ 8.124	+ 9.552	+ 9.633
Kaolinite $\frac{1}{2} (\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4)$	+ 5.561	+ 5.561	+ 3.514	+ 5.228	---	+ 3.802	+ 3.714	+ 2.853	+ 3.705	---	+ 3.705

The solubility product of corundum is written as follows:


$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 = 9.633 - 1.5 \log [\text{H}_2\text{O}] \quad (15)$$

When  $\text{Al}_2\text{O}_3$  is involved in an Al-hematite solid solution ( $0.5(\text{Al}_y\text{Fe}_{1-y})_2\text{O}_3$ )  $y$ , being the corundum mole fraction, and  $\lambda_y$ , the corresponding activity coefficient. One has:

$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 = 9.633 + \log y + \log \lambda_y - 1.5 \log [\text{H}_2\text{O}] \quad (16)$$

If the solid solution is ideal,  $\lambda_y = 1$ .

The equilibrium reaction of kaolinite with pure diasporite yields:



with

$$\log [\text{SiO}_2] = 3.705 - 8.949 - \frac{1}{2} \log [\text{H}_2\text{O}] = 5.244 - \frac{1}{2} \log [\text{H}_2\text{O}] \quad (17)$$

When the activity of water decreases,  $\log [\text{SiO}_2]$  corresponding to the equilibrium diasporite-kaolinite increases while  $\log [\text{SiO}_2]$  corresponding to the equilibrium gibbsite-kaolinite decreases. When  $\log [\text{H}_2\text{O}] = -0.744$ , gibbsite and diasporite are in equilibrium, so that  $\log [\text{SiO}_2] = -4.87$  in both cases. An increase in the water activity favors gibbsite on the one hand and acts to the detriment of diasporite or boehmite on the other. The equilibrium conditions, when ideal Al-goethite ( $\lambda = 1$ ) and kaolinite coexist together, are those given by equalizing the expression  $\log [\text{Al}^{3+}]/[\text{H}^+]^3$  in eqs (9) and (14):

$$\begin{aligned} \log [\text{Al}^{3+}]/[\text{H}^+]^3 \text{ in aqueous solution} &= 3.705 - \log [\text{SiO}_2] - 2.5 \log [\text{H}_2\text{O}] \\ &= 8.949 + \log x - 2 \log [\text{H}_2\text{O}] \end{aligned}$$

So that:

$$\log x = -5.244 - \log [\text{SiO}_2] - 0.5 \log [\text{H}_2\text{O}] \quad (18)$$

When the water activity is equal to one, and at the equilibrium with kaolinite and quartz ( $\log [\text{SiO}_2] = -4$ ), the  $\text{AlO}(\text{OH})$  mole fraction is calculated as follows:

$$\log x = -5.244 + 4 = -1.244$$

$$x = 0.06, \text{ corresponding to } (\text{Al}_{0.06}\text{Fe}_{0.94})\text{O}(\text{OH})$$

If we suppose that the equilibrium between Al-goethite, kaolinite, and quartz is realized in a dry environment ( $[\text{H}_2\text{O}] = 0.8$ ;  $-0.5 \log [\text{H}_2\text{O}] = -0.194$ )

$$\log x = -5.244 + 4 + 0.194 = -1.05$$

$$x = 0.08$$

It corresponds to  $(\text{Al}_{0.08}\text{Fe}_{0.92})\text{O}(\text{OH})$ . As the activity of water or the activity of silica decreases, Al-content in goethite increases.

For a given water activity,  $x$  increases as the activity of silica decreases; for a given activity of silica,  $x$  increases as the water activity decreases.

The equilibrium conditions when Al-hematite and kaolinite coexist are written, according to eqs (9) and (16), as follows:

$$\begin{aligned}\log [\text{Al}^{3+}]/[\text{H}^+]^3 \text{ in aqueous solution} &= 3.705 - \log [\text{SiO}_2] - 2.5 \log [\text{H}_2\text{O}] \\ &= 9.633 + \log y - 1.5 \log [\text{H}_2\text{O}]\end{aligned}$$

Then

$$\log [\text{SiO}_2] + \log y + \log [\text{H}_2\text{O}] = -5.928$$

For water activity equal to the unity ( $\log [\text{H}_2\text{O}] = 0$ ) and if one supposes that quartz is controlling the silica activity ( $\log [\text{SiO}_2] = -4$ ) then:

$$\log y = -1.928$$

$$y = 0.01$$


Hematite is almost pure:  $(\text{Al}_{0.01} \text{Fe}_{0.99})_2 \text{O}_3$ .

If activities of water and silica decrease together, and if for example,  $\log [\text{H}_2\text{O}] = -0.14$  (equilibrium gibbsite-boehmite) and if  $\log [\text{SiO}_2] = -4.87$  (equilibrium kaolinite-gibbsite-boehmite) then,  $\log y$  becomes:


$$\log y = -0.918 \text{ corresponding to: } (\text{Al}_{0.12} \text{Fe}_{0.88})_2 \text{O}_3.$$

When the silica and water activities decrease, the amount of aluminum in hematite increases. Furthermore, when silica activity becomes too low, kaolinite is no longer stable and the amounts of aluminum in goethite and hematite are only controlled by water activity.


*E. Gibbsite, boehmite, Al-goethite, and Al-hematite equilibrium conditions.*—The equilibrium reactions between gibbsite, boehmite, and diaspore yield the following conditions. When boehmite forms:



When the activity of water is sufficiently low, boehmite, instead of gibbsite, is supposed to be formed. If the water activity diminishes further, corundum forms:



Corundum may form in very dry environments ( $a_w < 0.04$ ). However, such a very low value is probably never reached in soils, and consequently corundum does not appear as a low temperature mineral. Furthermore, diaspore appears as less stable than boehmite. The transformation of gibbsite into diaspore involves the following conditions:



However, if diaspora appears as an endmember taking part in an Al-goethite ideal solid solution, we may have at equilibrium with gibbsite (eqs 3 and 14):

$$\begin{aligned}\log [\text{Al}^{3+}]/[\text{H}^+]^3 \text{ in aqueous solution} &= 8.949 + \log x - 2 \log [\text{H}_2\text{O}] \\ &= 8.205 - 3 \log [\text{H}_2\text{O}]\end{aligned}$$

So that, at equilibrium:

$$\log x = -0.744 - \log [\text{H}_2\text{O}]$$

In humid conditions, water activity is equal to unity ( $\log [\text{H}_2\text{O}] = 0$ ). The mole fraction of diaspora in Al-goethite at equilibrium with gibbsite is fixed:

$$\log x = -0.744$$

$$x = 0.18, \text{ corresponding to } (\text{Al}_{0.18} \text{Fe}_{0.82})\text{O}(\text{OH})$$

In dryer conditions, when gibbsite transforms to boehmite, the activity of water is lower ( $\log [\text{H}_2\text{O}] = -0.14$ ). Then, the mole fraction of diaspora in goethite is also fixed. It is, therefore, not dependent upon the water activity:

$$\log x = \log K_{\text{sp}} \text{ boehmite} - K_{\text{sp}} \text{ diaspora}$$

$$\log x = -0.604$$

$$x = 0.25, \text{ corresponding to } (\text{Al}_{0.25} \text{Fe}_{0.75})\text{O}(\text{OH})$$

When at equilibrium with boehmite, the water activity decreases, the amounts of aluminum in goethite or hematite also increase.

Four situations are represented in the equilibrium diagrams of figure 6. When iron is not represented in the system, the three straight lines delineate a field corresponding to an undersaturated solution in which the three minerals gibbsite, kaolinite, and quartz are unstable. The positions of the equilibrium straight lines of gibbsite and kaolinite depend only on the activity of water (fig. 6A and B). When iron is present in the system, Al-goethite (fig. 6C) or Al-hematite (fig. 6D) may form. These two minerals fall in the stability field occupied by undersaturated solutions. For a given activity of water the amount of substitution of the diaspora component in Al-goethite or the corundum component in Al-hematite both diminish when  $\log [\text{Al}^{3+}]/[\text{H}^+]^3$  decreases (fig. 6C and D).

*F. Al-goethite and Al-hematite equilibrium conditions.*—In the case of ideal solid solutions, the equilibrium conditions when Al-goethite and Al-hematite are together in equilibrium are written as follows. For  $(\text{Al}_x \text{Fe}_{1-x})\text{O}(\text{OH})$  and  $\text{Al}_y \text{Fe}_{1-y}\text{O}_{0.15}$ , the two partial equilibrium reactions are (eqs 1, 2, 14, and 16):

$$\begin{aligned}\log [\text{Al}^{3+}]/[\text{H}^+]^3 &= \log K_{\text{sp}} \text{ diaspora} - 2 \log [\text{H}_2\text{O}] + \log x \\ &= \frac{1}{2} \log K_{\text{sp}} \text{ corundum} - \frac{3}{2} \log [\text{H}_2\text{O}] + \log y\end{aligned}$$

together with (eqs 5 and 6):






Fig. 6. Equilibrium diagrams of kaolinite, gibbsite, Al-goethite, and Al-hematite in four different cases: (A)  $a_w = 1$  and no iron in the system; (B)  $a_w = 0.5$  and no iron in the system; (C)  $a_w = 1$  and Al-goethite forms; (D)  $a_w = 0.5$  and Al-hematite forms. The tie-lines represent the mole fractions of the diaspore component in Al-goethite (C) or the mole fraction of the corundum component in Al-hematite (D).

$$\begin{aligned}\log [\text{Fe}^{3+}]/[\text{H}^+]^3 &= \log K_{\text{sp}} \text{ goethite} - 2 \log [\text{H}_2\text{O}] + \log (1-x) \\ &= \frac{1}{2} K_{\text{sp}} \text{ hematite} - \frac{3}{2} \log [\text{H}_2\text{O}] + \log (1-y)\end{aligned}$$

$x$  and  $y$  are functions of the water activity

$$\log x/y = \frac{1}{2} \log K_{\text{sp}} \text{ hematite} - \log K_{\text{sp}} \text{ goethite} + \frac{1}{2} \log [\text{H}_2\text{O}]$$

and

$$\log (1-x)/(1-y) = \log K_{\text{sp}} \text{ diaspore} - \frac{1}{2} \log K_{\text{sp}} \text{ corundum} + \frac{1}{2} \log [\text{H}_2\text{O}]$$

When the water activity diminishes, the mole fractions of goethite and diaspore diminish:  $x$  and  $y$  are determined by solving simultaneously the two equations. The amounts of Al-goethite and Al-hematite depend on the amounts of Al and Fe in the system.

For pure diaspore and corundum:  $x = 1$  and  $y = 1$ , the equilibrium condition yields, as already shown

$$\begin{aligned}\frac{1}{2} \log [\text{H}_2\text{O}] &= \log K_{\text{sp}} \text{ diaspore} - \frac{1}{2} \log K_{\text{sp}} \text{ corundum} \\ &= 8.949 - 9.633 = -0.684 \\ [\text{H}_2\text{O}] &= 0.21\end{aligned}$$

For pure goethite and hematite:  $x = 0$  and  $y = 0$ . The equilibrium condition is the following

$$\begin{aligned}\frac{1}{2} \log [\text{H}_2\text{O}] &= \log K_{\text{sp}} \text{ goethite} - \frac{1}{2} \log K_{\text{sp}} \text{ hematite} \\ &= +0.819 - 0.846 = -0.027 \\ [\text{H}_2\text{O}] &= 0.88\end{aligned}$$

The dehydration of goethite into hematite appears at higher water activities than for the dehydration of diaspore into corundum. A decrease in water activity, when the two solid solutions are in equilibrium, must induce an increase of the aluminum content in Al-hematite. Furthermore, in contradiction to the model proposed by Yapp (1983), it is evident that the two solid solutions in equilibrium cannot show the same composition. The substitution rates of Fe by Al are obviously different in the two minerals and are changing as functions of the water activity and the composition of the system.

*G. Fe<sup>3+</sup>-kaolinite.*—The solubility product equations of kaolinite and Fe<sup>3+</sup>-kaolinite are written as follows

$$\begin{aligned}\frac{1}{2} \text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4 + 3 \text{H}^+ &\rightleftharpoons \text{Al}^{3+} + \text{SiO}_{2(\text{aq})} + \frac{5}{2} \text{H}_2\text{O} \\ \log [\text{Al}^{3+}]/[\text{H}^+]^3 + 2 \log [\text{SiO}_2] + 2.5 \log [\text{H}_2\text{O}] &= 3.705\end{aligned}$$

and

$$\begin{aligned}\frac{1}{2} \text{Fe}_2\text{Si}_2\text{O}_5(\text{OH})_4 + 3 \text{H}^+ &\rightleftharpoons \text{Fe}^{3+} + \text{SiO}_{2(\text{aq})} + \frac{5}{2} \text{H}_2\text{O} \\ \log [\text{Fe}^{3+}]/[\text{H}^+]^3 + 2 \log [\text{SiO}_2] + 2.5 \log [\text{H}_2\text{O}] &= -3.027\end{aligned}$$

The solubility of Fe<sup>3+</sup>-kaolinite is derived from Tardy (1982). It may be calculated in order to satisfy the equilibrium between a 1 percent Fe<sup>3+</sup>-kaolinite in equilibrium with a 20 percent Al-goethite. This is what is observed in lithomarges where kaolinite and goethite coexist. For a Fe<sup>3+</sup>-kaolinite solid solution ((Fe<sub>z</sub><sup>3+</sup> Al<sub>1-z</sub><sup>3+</sup>) Si<sub>2</sub>O<sub>5</sub>(OH)<sub>4</sub>) one obtains:

$$\log [\text{Fe}^{3+}]/[\text{Al}^{3+}] = 1/2 \log K_{\text{sp}} \text{Fe}^{3+}\text{-kaolinite} - 1/2 \log K_{\text{sp}} \text{kaolinite} \\ + \log (z/(1-z))$$

one finds:

$$\log [\text{Fe}^{3+}]/[\text{Al}^{3+}] = -1.823 - 3.705 + \log (z/(1-z)) \\ = -5.528 + \log (z/(1-z))$$

At the equilibrium with an Al-goethite solid solution for which  $x = 0.2$ ,  $(1-x) = 0.8$ , one has:

$$\log [\text{Fe}^{3+}]/[\text{Al}^{3+}] = 0.819 - 8.949 + \log (x/(1-x)) \\ = -8.130 + 0.602 = -7.528$$


and

$$\log (z/(1-z)) = -7.528 + 5.528 = -2$$

In that case:

$$z = 0.01$$

This corresponds to the Fe<sup>3+</sup>-kaolinite formula:



At equilibrium with an Al-goethite of a given composition, the amount of Fe<sup>3+</sup> in kaolinite should increase when the kaolinite crystallinity diminishes, that is, when the kaolinite solubility increases.

The distribution of Al and Fe in kaolinite, goethite, and hematite has been calculated by the use of a computer program called MINEQ. This program is based on the minimization of the Gibbs free energies of the systems of a given composition and is discussed in the next section.

#### AN IDEAL SOLID SOLUTION MODEL DISTRIBUTION OF PHASES BY THE GIBBS FREE ENERGY MINIMIZATION

On the basis of equilibrium, the element distribution among phases and solid solutions can be calculated by using a method of minimization of Gibbs free energy. A corresponding computer program called MINEQ was designed by Perret (1983), Perret, Fritz, and Tardy (1984) and applied by Didier (1983) to the system SiO<sub>2</sub>-Al<sub>2</sub>O<sub>3</sub>-Fe<sub>2</sub>O<sub>3</sub>-H<sub>2</sub>O. The distribution of the elements among solid solutions is calculated on the basis of ideal mixing and as a function of element composition and water activity. All the data presented refer to 1 mole of Fe<sub>2</sub>O<sub>3</sub>. The amounts of Al<sub>2</sub>O<sub>3</sub> and SiO<sub>2</sub> change from one set to another. The amount of water is limited or unlimited depending on the capability of profiles to be either closed or open for circulating water. In the first case, water activity should be lower


than unity. In the second case, it can be unity (saturated regime) but also lower than unity (unsaturated regime).

The  $K_{sp}$  and the corresponding Gibbs free energies used in the model are those of table 4, column 11, and table 3, column 11, respectively.

*A. A sample calculation.*—A sample calculation, in the system  $\text{Fe}_2\text{O}_3$ – $\text{Al}_2\text{O}_3$ – $\text{H}_2\text{O}$  is given as follows. The molal composition

$$\begin{aligned}\text{Fe}_2\text{O}_3 &= 1 & (\text{Fe} = 2) \\ \text{Al}_2\text{O}_3 &= 0.1 & (\text{Al} = 0.2) \\ \text{SiO}_2 &= 0 \\ \text{H}_2\text{O} &= \infty \\ a_w &= 0.7\end{aligned}$$


can be distributed among an infinite number of mineral assemblages of various compositions, but only one assemblage and composition correspond to equilibrium. The calculated equilibrium distribution is the following:



The  $\Delta G$  of the system based on ideal solid solutions yields:

$$\begin{aligned}\Delta G_{\text{system}} &= 0.5582 \Delta G_f^\circ \text{Al-goethite} + 1.5775 \frac{1}{2} \text{Al-hematite} \\ &= -885.558 \text{ kJ/mole}\end{aligned}$$


Two other assemblages listed below are among many stoichiometrically equivalent. They are, for instance, either:



for which

$$\Delta G_{\text{system}} = -885.500 \text{ kJ/mole}$$

or:



for which:

$$\Delta G_{\text{system}} = -885.382 \text{ kJ/mole}$$

$\Delta G$  of the system is at a minimum only for the composition corresponding to equilibrium.

Furthermore, the elemental and mineral distributions satisfy the following equilibrium relations (from eqs 5 and 6):

$$\begin{aligned}\log [\text{Fe}^{3+}]/[\text{H}^+]^3 (\text{goethite}) &= \log [\text{Fe}^{3+}]/[\text{H}^+]^3 (\frac{1}{2} \text{hematite}) \\ &= \log X + \log K_{sp} \text{goethite} - 2 \log [\text{H}_2\text{O}] = 1.113 \\ &= \log Y + \log K_{sp} \frac{1}{2} \text{hematite} - 1.5 \log [\text{H}_2\text{O}] = 1.113\end{aligned}$$

Also, from eqs 1, 2, 13, and 15, the equilibrium conditions are satisfied:

$$\log [\text{Al}^{3+}]/[\text{H}^+]^3 (\text{diaspore}) = \log [\text{Al}^{3+}]/[\text{H}^+]^3 (\frac{1}{2} \text{corundum})$$

$$= \log (1-X) + \log K_{sp} \text{ diaspore} - 2 \log [\text{H}_2\text{O}] = 8.407$$

$$= \log (1-Y) + \log K_{sp} \frac{1}{2} \text{ corundum} - 1.5 \log [\text{H}_2\text{O}] = 8.407$$

*B. Data obtained in an excess of water.*—In table 5, the results are shown for an excess of water so that the activity is equal to unity. For one mole of  $\text{Fe}_2\text{O}_3$ , the number of moles of  $\text{Al}_2\text{O}_3$  changes from 0 to 1, and the number of moles of  $\text{SiO}_2$  varies from 0 to 0.25.

All the results presented are based on 1 mole of  $\text{Fe}_2\text{O}_3$ , that is, 2 atoms of Fe. If, for example, aluminum in the systems is given as  $\text{Al}_2\text{O}_3 = 0.1$ , the number of atoms of Al = 0.2, and the mole fractions are respectively equal to:

$$\text{Al} = \frac{0.2}{0.2 + 2} = 0.0909 \text{ and } \text{Fe} = \frac{2}{0.2 + 2} = 0.9091$$

It is seen that hematite never forms in such conditions. Furthermore, quartz appears when silica content increases and when the system is poor in alumina. If the aluminum content increases over 0.10, gibbsite is formed, and consequently the amount of aluminum in goethite remains constant ( $\text{Fe}_{0.82} \text{Al}_{0.18} \text{O}(\text{OH})$ ). If silica and alumina increase together, kaolinite forms. Furthermore, if for a given amount of alumina (0.25), silica increases, the amount of kaolinite increases, the amount of goethite formed increases, but the Al substitution rate in goethite decreases.

In table 6 the reported results are obtained also for an excess of water of activity lower than unity ( $a_w = 0.7$ ). A similar picture is reached. However for low aluminum and for high silica contents, hematite tends to form instead of goethite. Therefore, goethite or a goethite and hematite mixture inhibit gibbsite formation up to an  $\text{Al}_2\text{O}_3$  content equal to 0.5. Here, the quantities of gibbsite formed are lower than in the previous case. Furthermore, the substitution rate of goethite by diaspore is higher in water of low activity than in water of high activity.

One can note that the differences in free energy are very low, on the order of a few hundreds of Joules. This is the reason the calculations presented in tables 5 and 6 may suffer some discrepancies. The compositions calculated are not precise, and the calculation procedure may still be improved.

Some of these results are reported on figure 7 where for one mole of  $\text{Fe}_2\text{O}_3$ , the composition and the stability fields of the hematite and goethite solid solutions are reported as functions of the water activity and aluminum content. In such conditions of excess water, hematite does not appear as associated with gibbsite or boehmite. This is observed also when the amount of water is reduced.

*C. Results obtained when the amount of water is limited.*—If we suppose, for example, that the porosity of the material equal to 40 percent is closed and that this porosity is initially filled by water, then the hydrated mineral formation (gibbsite, goethite) is limited, and the stability field of hematite is considerably enhanced. However, hematite and gibbsite do not coexist. If the activity of water decreases and if the water con-

TABLE 5

Composition and quantity of Al-goethite and Al-hematite associated with gibbsite, kaolinite, and quartz for various  $\text{SiO}_2$  or  $\text{Al}_2\text{O}_3$  content (1 mole of  $\text{Fe}_2\text{O}_3$ ,  $a_w = 1$  and excess of water)

$\text{Al}_2\text{O}_3$	0.00	0.05	0.10	0.25	0.50	1.00
$\text{SiO}_2$						
0	2 FeOOH	2.10 ( $\text{Fe}_{0.95}\text{Al}_{0.05}$ )OOH	2.20 ( $\text{Fe}_{0.91}\text{Al}_{0.09}$ )OOH	2.47 ( $\text{Fe}_{0.81}\text{Al}_{0.19}$ )OOH 0.03 G	2.47 ( $\text{Fe}_{0.81}\text{Al}_{0.19}$ )OOH 0.53 G	2.47 ( $\text{Fe}_{0.81}\text{Al}_{0.19}$ )OOH
0.05	2 FeOOH 0.05 Q	2.10 ( $\text{Fe}_{0.95}\text{Al}_{0.05}$ )OOH 0.05 Q	2.15 ( $\text{Fe}_{0.93}\text{Al}_{0.07}$ )OOH 0.03 K	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.03 K	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.03 K 0.50 G	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.03 K 1.50 G
0.10	2 FeOOH 0.10 Q	2.10 ( $\text{Fe}_{0.95}\text{Al}_{0.05}$ )OOH 0.10 Q	2.13 ( $\text{Fe}_{0.94}\text{Al}_{0.06}$ )OOH 0.02 Q	2.41 ( $\text{Fe}_{0.83}\text{Al}_{0.17}$ )OOH 0.05 K 0.04 K	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.05 K 0.46 G	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.05 K 1.46 G
0.25	2 FeOOH 0.25 Q	2.10 ( $\text{Fe}_{0.95}\text{Al}_{0.05}$ )OOH 0.25 Q	2.13 ( $\text{Fe}_{0.94}\text{Al}_{0.06}$ )OOH 0.17 Q 0.04 K	2.25 ( $\text{Fe}_{0.89}\text{Al}_{0.11}$ )OOH 0.13 K	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.13 K 0.30 G	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.13 K 1.30 G

TABLE 6

Composition and quantity of Al-goethite and Al-hematite associated with gibbsite, quartz, and kaolinite for various  $\text{SiO}_2$  or  $\text{Al}_2\text{O}_3$  content (1 mole of  $\text{Fe}_2\text{O}_3$ ,  $a_w = 0.70$  and excess of water)

$\text{Al}_2\text{O}_3$	0.00	0.05	0.10	0.25	0.50	1.00
$\text{SiO}_2$						
0	2 $\text{FeO}_{1.5}$	0.37 ( $\text{Fe}_{0.87}\text{Al}_{0.13}$ )OOH 1.73 ( $\text{Fe}_{0.97}\text{Al}_{0.03}$ )O.	1.34 ( $\text{Fe}_{0.87}\text{Al}_{0.13}$ )OOH 0.86 ( $\text{Fe}_{0.97}\text{Al}_{0.03}$ )O <sub>1.5</sub>	2.50 ( $\text{Fe}_{0.80}\text{Al}_{0.20}$ )OOH	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 0.30 G	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 1.30 G
0.05	2 $\text{FeO}_{1.5}$ 0.05 Q	2.04 ( $\text{Fe}_{0.98}\text{Al}_{0.02}$ )O <sub>1.5</sub> 0.03 K	0.86 ( $\text{Fe}_{0.87}\text{Al}_{0.13}$ )OOH 0.64 ( $\text{Fe}_{0.97}\text{Al}_{0.03}$ )O <sub>1.5</sub> 0.03 K	2.44 ( $\text{Fe}_{0.82}\text{Al}_{0.18}$ )OOH 0.03 K	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 0.03 K 0.25 G	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 0.03 K 1.25 G
0.10	2 $\text{FeO}_{1.5}$ 0.10 Q	2.04 ( $\text{Fe}_{0.98}\text{Al}_{0.02}$ )O <sub>1.5</sub> 0.03 K 0.04 Q	2.15 ( $\text{Fe}_{0.93}\text{Al}_{0.07}$ )OOH 0.03 K 0.04 Q	2.41 ( $\text{Fe}_{0.83}\text{Al}_{0.17}$ )OOH 0.05 K	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 0.05 K 0.20 G	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 0.05 K 1.20 G
0.25	2 $\text{FeO}_{1.5}$ 0.25 Q	2.04 ( $\text{Fe}_{0.98}\text{Al}_{0.02}$ )O <sub>1.5</sub> 0.03 K 0.19 Q	2.15 ( $\text{Fe}_{0.93}\text{Al}_{0.07}$ )OOH 0.03 K 0.19 Q	2.25 ( $\text{Fe}_{0.89}\text{Al}_{0.11}$ )OOH 0.13 K	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 0.13 K 0.04 G	2.70 ( $\text{Fe}_{0.74}\text{Al}_{0.26}$ )OOH 0.13 K 1.04 G




Fig. 7. Stability field of Al-goethite and Al-hematite as a function of water activity and  $\text{Al}_2\text{O}_3$  content in the system  $\text{Al}_2\text{O}_3\text{--Fe}_2\text{O}_3\text{--H}_2\text{O}$ . Tie-lines represent the mole fraction of diaspore in goethite or corundum in hematite.

tent diminishes, gibbsite is no longer stable with respect to boehmite, and goethite no longer stable with respect to hematite. Furthermore, for relatively low  $\text{Al}_2\text{O}_3$  contents, Al-goethite or Al-hematite can be more stable than gibbsite or boehmite, so that no aluminum phase can crystallize. The model confirms also that, when kaolinite is present together with goethite or hematite, the amount of aluminum substituted in the oxide or the oxyhydroxide decreases when the silica activity increases. The paragenesis indicates that the only free aluminum phase associated with hematite should be boehmite. When gibbsite and boehmite are excluded, goethite and hematite coexist; their aluminum content increases with the total aluminum content in the system; therefore, the substitution rate of corundum in hematite is always lower than the substitution rate of diaspore in goethite.

#### CONCLUDING REMARKS

The question of the equilibrium conditions in soils, as functions of water activity and the amount of available water, has been discussed by Chesworth (1972, 1975) for the system  $\text{SiO}_2\text{--Al}_2\text{O}_3\text{--Fe}_2\text{O}_3\text{--H}_2\text{O}$  with emphasis on gibbsite and boehmite stability. The gibbsite-kaolinite equilib-



rium has been analyzed by Mattigod and Kittrick (1980), Pédro and Delmas (1980), Sposito (1981), and Tardy (1982). The diminution of the water activity due to the transition from the saturated regime to the unsaturated one is clearly realized to be an important factor, which can be responsible for phase boundary displacement.

From careful petrographic observations made on the distribution of goethite and hematite in ferricretes and lateritic red soils, conclusions may be drawn.

1. Goethite commonly forms in young bauxitic profiles and is preserved in the lower parts of the old bauxitic formations. Here, goethite and gibbsite coexist in a very common lateritic paragenesis. Furthermore, when, in the old bauxites, boehmite can take the place of gibbsite, hematite forms instead of goethite.

2. Goethite is also the dominant ferric mineral in the permanently humid tropical red or yellow lateritic soils in which kaolinite dominates. Here, the typical paragenesis is quartz, kaolinite, and goethite. The iron oxihydroxide is intimately associated with kaolinite, forming small and tiny *grumeaux*. In that case, goethite does not replace kaolinite but "coexists with it."

3. In seasonally varying tropical soils, goethite predominates only in downslope profiles (wet, most of the time) in the mottle zones, where the water table fluctuates and in saprolites and lithomarges below the water table. Here again, goethite and kaolinite coexist with no replacement of the clay by the oxihydroxide.

4. Goethites of these types are all more or less aluminous and may contain up to 30 mole percent diaspore. The aluminum content is controlled by the aluminum available in solution in which goethites form. As shown for example by Fritz and Tardy (1974, 1976) or Fritz (1981), aluminum in solution increases during a mineral sequence, as one goes from montmorillonite to kaolinite plus quartz, then to kaolinite without quartz, and finally to gibbsite. From dry to humid climates, from downslopes to upslopes, from bottom to the top of the profiles, drainage and leaching increase, silica concentrations in solutions decrease, while aluminum in solution and in goethite increase together. The aluminum content of goethites coexisting with kaolinite is controlled by the silica activity in solution, probably much more than by other factors that accompany the leaching such as pH, organic matter content, *et cetera*.

5. Hematite appears at the top of some old bauxitic profiles, together with boehmite and sometimes with diaspore. Generally, hematite is not abundant when gibbsite dominates.

6. Therefore, under seasonally varying tropical climates, in the upper parts of the upslope profiles above the mottled zones, in concretioned horizons, hematite appears as a very common mineral. Then, the formation of hematite in concretions follows the formation of goethite in mottles.

7. When a concretion develops, iron is enriched in zones of small size porosity, in preference to the areas in which large sized pores pre-

dominate. This is the reason hematite forms predominantly in microenvironments in which kaolinite was initially abundant. Therefore, instead of merely coexisting, as observed for goethite, hematite replaces kaolinite and becomes aluminous. The reason for this replacement is essentially the instability of kaolinite in the unsaturated zone. Furthermore, the hematite is extremely finely crystallized.

8. In the upper parts of ferricretes, dismantling takes place, and large channels develop. Along the margins of the oxide concretions and in contact with large size pores periodically visited by percolating waters, aluminous hematites alter into secondary aluminum-rich goethites. These secondary minerals are transformed, in turn, into less and less aluminous goethites, relatively better crystallized than the replaced hematites.

9. At sufficiently high activities of water and at sufficiently low activities of silica in solution, gibbsite forms together with Al-goethite. The aluminum content in goethite is fixed, when the activity of water is fixed. If the water activity diminishes or if the activity of silica increases if the ratio of Al/Fe diminishes, gibbsite does not form. Thus, the stability field of gibbsite is limited by either boehmite, kaolinite, or Al-goethite formation. Hematite forms when the activity of water and the ratio Al/Fe are both low. When Al-hematite or Al-goethite form alone, their aluminum content is determined by the composition of the system; when they form together, their compositions are determined only by the activity of water. In this case, the amount of each mineral is then determined by the activity of water and by the Al/Fe ratio in the system.

10. Finally, the parameters that control the relative stability of hematites and goethites are: (1) grain size of particles, (2) aluminum, iron, and water content in the system, (3) activity of silica when kaolinite or smectite is present, (4) activity of water, that is, saturated or non-saturated regime and in the latter case pore size in which minerals form.

#### REFERENCES

- Angel, B. R., and Hall, P. L., 1973, Electron spin resonance studies of kaolins: Internat. Clay Conf., Madrid, Proc., p. 47-60.
- Bardossy, G., 1968, *Le gisement de bauxite d'Eplény*: Budapest, Földtani Közlöny, v. 98, p. 408-442.
- 1982, *Karst bauxites. Bauxites deposits on carbonate rocks*: New York, Elsevier, 441 p.
- Beneslavsky, S. I., 1957, New aluminum-bearing minerals in bauxites: Akad. Nauk SSSR, Doklady, v. 113, p. 367-369.
- Berner, R. A., 1969, Goethite stability and the origin of red beds: *Geochim. et Cosmochim. Acta*, v. 33, p. 267-273.
- 1971, *Principles of chemical sedimentology*: New York, McGraw-Hill, 240 p.
- Bischoff, J. L., 1969, Goethite-hematite stability relations with relevance to sea water and the Red Sea Brines system, in Degens, E. T., and Ross, D. A., eds., *Hot brines and recent heavy metal deposits in the Red Sea*: New York, Springer-Verlag Inc., p. 402-406.
- Boesman, E., and Schoemaker, D., 1961, Résonance paramagnétique de l'ion  $Fe^{3+}$  dans la kaolinite: *Acad. Sci. Paris Comptes rendus*, v. 252, p. 1931-1933.
- Boulangé, B., 1983, *Les formations bauxitiques latéritiques de Côte d'Ivoire*: Thèse, Univ. Paris VII, 341 p.
- Boulet, R., 1974, *Toposéquences de sols tropicaux en Haute-Volta. Equilibre et déséquilibre pédoclimatique*: Mém. Office de la Recherche Sci. et Tech. Outre-Mer (ORSTOM), Paris, 1978, 272 p.

- Bourrié, G., and Pédro, G., 1979, la notion de pF, sa signification physico-chimique et ses implications pédogénétiques. I. Signification physico-chimique. Relations entre pF et activité de l'eau: *Sci. Sol.*, v. 4, p. 313-322.
- Brunauer, S., Deming, L. S., Deming, W. E., and Teller, E., 1940, On a theory of Van der Waals adsorption of gases: *Am. Chem. Soc. Jour.*, v. 62, p. 1723-1732.
- Brunauer, S., and Emett, P. H., 1935, The use of Van der Waals adsorption isotherms in determining the surface area of iron synthetic ammonia catalysts. *Am. Chem. Soc. Jour.*, v. 57, p. 1734-1755.
- 1937, The use of Van der Waals adsorption isotherms in determining the surface area of various adsorbents. *Am. Chem. Soc. Jour.*, v. 59, p. 2682-2689.
- Brunauer, S., Emett, P. H., and Teller, E., 1938, Adsorption of gases in multimolecular layers: *Am. Chem. Soc. Jour.*, v. 60, p. 309-319.
- Caillère, S., Gatineau, L., and Hénin, S., 1960, Préparation à basse température d'hématite alumineuse: *Acad. Sci. Paris Comptes rendus*, v. 250, p. 3677-3679.
- Cantinolle, P., Didier, P., Meunier, J. D., Parron, C., Guendon, J. L., Bocquier, G., and Nahon, D., 1983, Kaolinites ferrifères et oxyhydroxydes de fer et d'alumine dans les bauxites des Canonettes (S.E. de la France): *Clay Minerals*, in press.
- Cases, J. M., Liétard, O., Yvon, Y., and Delon, J. F., 1981, Etude des propriétés cristallographiques et superficielles des kaolinites désordonnées: *Bull. Minéralogie, supp.*, v. 104, 5, p. 55.
- Chesworth, W., 1972, The stability of gibbsite and boehmite at the surface of the earth: *Clays and Clay Minerals*, v. 20, p. 369-374.
- 1975, The system  $\text{SiO}_2\text{-AlO(OH)-Fe}_2\text{O}_3\text{-H}_2\text{O}$  and the kaolinitic stage of the goethite facies: *Clays and Clay Minerals*, v. 23, p. 389-392.
- Chetelat, E. de, 1938, Le modelé latéritique de l'Ouest de la Guinée française: *Rev. Géographie Physique et Géologie Dynamique*, v. 11, 1, 119 p.
- Combes, P. J., 1969, Recherches sur la genèse des bauxites dans le Nord-Est de l'Espagne, le Languedoc et l'Ariège (France): *Mém. Centre Recherche Géologie Hydrogéologie*, Montpellier, 342 p.
- Correns, C. W., and Von Engelhardt, W., 1941, Röntzenographische Untersuchungen über den Mineralbest und Sedimentärer Eisenerze: *Nachr. Akad. Wiss. Göttingen, Math.-Phys. Kl.*, v. 2, p. 131-137.
- Dandurand, J. L., Mizele, J., Schott, J., Bourgeat, F., Valles, V., and Tardy, Y., 1982, Premiers résultats sur la solubilité de la silice amorphe dans les pores de petite taille. Variations du coefficient d'activité de la silice en fonction de l'activité de l'eau: *Sci. Géol. Bull.*, v. 35, 1-2, p. 71-79.
- Davey, B. G., Russel, J. D., and Wilson, M. J., 1975, Iron oxide and clay minerals and their relation to colors of red and yellow podzolic soils near Sydney, Australia: *Geoderma*, v. 14, p. 125-138.
- Delon, J. F., and Dellyes, R., 1967, Calcul du spectre de porosité des minéraux phyliteux: *Acad. Sci. Paris Comptes rendus*, v. 265, p. 1661-1664.
- D'Hoore, J., 1954, Etude statistique des teneurs en  $\text{Fe}_2\text{O}_3$  libre sur limon et argile pour les sols des trois régions naturelles du Congo Belge: *Internat. Conf. African Soils*, 2, Léopoldville, v. 14, 15 p.
- Didier, Ph., 1983, Paragenèses à oxydes et hydroxydes de fer dans les bauxites et les cuirasses ferrugineuses: Thèse 3ème cycle, Univ. Poitiers, 150 p.
- Didier, Ph., Fritz, B., Nahon, D., and Tardy, Y., 1983,  $\text{Fe}^{3+}$ -kaolinites, Al-goethites and Al-hematites in tropical ferricretes, in *Petrology of weathering and soils*, Nahon, D. and Noack, Y., eds.: *Sci. Géol. Mém.*, v. 71, p. 35-44.
- Didier, Ph., Perret, D., Tardy, Y., and Nahon, D., 1984, Equilibres entre kaolinites ferrifères, goethites alumineuses et hématites alumineuses dans les systèmes cuirassés. Rôle de l'activité de l'eau et de la taille des pores: *Sci. Géol. Bull.*, in press.
- Endredy, A. S., de, 1963, Estimation of free iron oxides in soils and clay by a photolytic method: *Clay Minerals Bull.*, v. 5, p. 209-217.
- Fayolle, M., 1979, Caractérisation analytique d'un profil d'argile à silex de l'Ouest du bassin de Paris: Thèse, 3ème cycle, Univ. Paris VII, 162 p.
- Fischer, A. G., 1963, Easy review of descriptive paleoclimatology. *Am. Jour. Sci.*, v. 261, p. 282-293.
- Fitzpatrick, R. W., and Schwertmann, U., 1982, Al-substituted goethite. An indicator of pedogenetic and other weathering environments in South Africa: *Geoderma*, v. 27, p. 335-347.
- Fölster, H., Noshrefi, N., and Ojenuga, A. G., 1971, Ferralitic pedogenesis on metamorphic rock, S.W. Nigeria: *Pedologie*, v. 20, 1, p. 95-124.

- Fritz, B., 1981, Etude thermodynamique et modélisation des réactions hydrothermales et diagénétiques: *Sci. Géol. Mém.*, v. 65, 197 p.
- Fritz, B., and Tardy, Y., 1974, Etude thermodynamique du système gibbsite, quartz, kaolinite, gaz carbonique. Application à la genèse des podzols et des bauxites. *Sci. Géol. Bull.*, v. 26, 4, 339-367.
- , 1976, Prediction of mineralogical sequences in tropical soils by a theoretical dissolution model: *Internat. Symposium on Water Rock Interaction*, Prague, Sept. 1974, p. 409-416.
- Garrels, R. M., and Christ, C. L., 1965, *Solutions, Minerals and Equilibria*: New York, Harper and Row, 450 p.
- Goldich, S. S., and Berquist, H. R., 1948, Aluminous lateritic soil of the republic of Haiti: *U.S. Geol. Survey Bull.* 954-C, p. 63-111.
- Hall, P. L., 1980, The application of electron spin resonance spectroscopy to studies of clay minerals: I. Isomorphous substitutions and external surface properties: *Clay Minerals*, v. 15, p. 321-335.
- Helgeson, H. C., Delany, J. M., Nesbitt, H. W., and Bird, D. K., 1978, Summary and critique of the thermodynamic properties of rock-forming minerals: *Am. Jour. Sci.*, v. 278-A, 229 p.
- Hem, J. D., 1970, Aluminum. 13-G. Behavior during weathering and alteration of rocks: *Handbook of Geochemistry*, v. 2, no. 2, p. 13G1-13L3.
- Hem, J. D., and Lind, J., 1974, Kaolinite synthesis at 25°C: *Science*, v. 184, p. 1171-1173.
- Hem, J. D., and Roberson, C. E., 1967, Form and stability of aluminum hydroxide complexes in dilute solution: *U.S. Geol. Survey Water Supply Paper* 1827 A, 55 p.
- Hem, J. D., Roberson, C. E., Lind, C. J., and Polzer, W. L., 1973, Chemical interactions of aluminum with aqueous silica at 25°C: *U.S. Geol. Survey Water Supply Paper* 1827 E, 57 p.
- Hemingway, B. S., and Robie, R. A., 1977a, The entropy and Gibbs free energy of formation of the aluminum ion: *Geochim. et Cosmochim. Acta*, v. 41, p. 1402-1404.
- , 1977b, Enthalpies of formation of low albite ( $\text{NaAlSi}_3\text{O}_8$ ), gibbsite ( $\text{Al}(\text{OH})_3$ ), and  $\text{NaAlO}_2$ ; revised values for  $\Delta H^\circ_{f,298}$  and  $\Delta G^\circ_{f,298}$  of some aluminosilicate minerals: *U.S. Geol. Survey Jour. Research*, v. 5, p. 413-429.
- Hemingway, B. S., Robie, R. A., and Kittrick, J. A., 1978, Revised values for the Gibbs free energy of formation of  $[\text{Al}(\text{OH})_4]_{\text{aq.}}^-$ , diaspore, boehmite and bayerite at 298.15 K and 1 bar, the thermodynamic properties of kaolinite to 800 K and 1 bar, and the heats of solution of several gibbsite samples: *Geochim. et Cosmochim. Acta*, v. 42, p. 1533-1543.
- Herbillon, A. J., Mestdagh, M. M., Vielvoye, L., and Derouane, E. G., 1976, Iron in kaolinite with special references to kaolinite from tropical soils: *Clay Minerals*, v. 11, 3, p. 201-220.
- Hillel, D., 1971, *Soils and water*: New York, Academic Press, 288 p.
- Iler, R. K., 1979, *The chemistry of silica*: Wiley-Interscience, 866 p.
- Janot, C., and Gibert, H., 1970, Les constituants du fer dans certaines bauxites naturelles étudiées par effet Mössbauer: *Soc. française Minéralogie Cristallographie Bull.*, v. 90, p. 213-223.
- Janot, C., Gibert, H., de Grammont, X., and Biais, R., 1971, Etude des substitutions Al-Fe dans des roches latéritiques. *Soc. française Minéralogie Cristallographie Bull.*, v. 94, p. 367-380.
- Janot, C., Gibert, H., and Tobias, C., 1973, Caractérisation de kaolinites ferrifères par spectrométrie Mössbauer. *Soc. française Minéralogie, Cristallographie Bull.*, v. 96, p. 281-291.
- Kämpf, N., and Schwertmann, U., 1983, Goethite and hematite in a climosequence in a southern Brazil and their application in classification of kaolinitic soils: *Geoderma*, v. 29, p. 27-39.
- Keenan, A. G., Mooney, R. W., and Wood, L. A., 1951, The relation between exchangeable ions and water adsorption on kaolinite: *Jour. Phys. Colloidal Chemistry*, v. 55, p. 1462-1474.
- Kehres, A., 1983, Isothermes de déshydratation des argiles. Energies d'hydratation. Diagrammes de pores. Surfaces internes: Thèse, Univ. Paul Sabatier, Toulouse, 163 p.
- Lamoureux, M., 1972, Etat et comportement du fer dans les sols formés sur roches carbonatées au Liban: *Sci. Sol*, v. 1, p. 85-101.
- Lamoureux, M., Loyer, J. Y., Bouleau, A., Janot, C., 1977, Formes du fer des sols rouges et bruns fersiallitiques. Application de la spectrométrie Mössbauer: *Cahier Office de la Recherche Sci. et Tech. Outre-Mer (ORSTOM)*, Paris, sér. Pédologie, v. 15, 2, p. 199-210.

- Langmuir, D., 1969, The Gibbs free energies of substances in the system Fe-O<sub>2</sub>-H<sub>2</sub>O-CO<sub>2</sub> at 25°C: U.S. Geol. Survey Prof. Paper 650 B, p. 180-184.
- 1971, Particle size effect on the reaction goethite = hematite + water: *Am. Jour. Sci.*, v. 271, p. 147-156.
- Leprun, J. C., 1971, Premières observations sur les toposéquences à amont cuirassé en Haute-Volta orientale: rôle de la pédogenèse dans la destruction des cuirasses et le façonnement du modelé: *Bull. Liaison, Thème B, Office de la Recherche Sci. et Tech. Outre-Mer (ORSTOM)*, Paris, v. 2, p. 39-53.
- 1972, Cuirasses ferrugineuses autochtones et modelé des bas reliefs des pays cristallins de Haute-Volta orientale. *Acad. Sci. Paris Comptes rendus*, v. 275 D, p. 1207-1210.
- 1979, Les cuirasses ferrugineuses des pays cristallins de l'Afrique Occidentale sèche. Genèse. Transformation. Dégradation: *Sci. Géol. Mém.* 58, 244 p.
- Maignien, R., 1958, Le cuirassement des sols en Guinée. *Afrique Occidentale: Mém. Service Carte Géol. Alsace-Lorraine*, v. 16, 239 p.
- Malden, P. J., and Meads, R. E., 1967, Substitution by iron in kaolinite: *Nature*, v. 215, p. 844-846.
- Mattigod, S. V., and Kittrick, J. A., 1980, Temperature and water activity as variables in soils mineral activity diagrams: *Soil Sci. Soc. America Proc.*, v. 44, p. 149-154.
- Mendelovici, E., Yariv, S. H., and Villalba, R., 1979, Iron bearing kaolinite in Venezuelan laterites. I. Infrared spectroscopy and chemical dissolution evidences: *Clay Minerals*, v. 14, 4, p. 323-331.
- Mestdagh, M. M., Vielvoye, L., and Herbillon, A. J., 1980, Iron in kaolinite: II. The relationship between kaolinite and iron content: *Clay Minerals*, v. 15, 1, p. 1-13.
- Michel, P., 1973, Les bassins des fleuves Sénégal et Gambie. Etude géomorphologique: *Mém. Office Recherche Sci. Tech. Outre-Mer (ORSTOM)*, Paris, v. 63, 1, 2, 3, 752 p.
- Millot, G., 1964, *Géologie des argiles*: Paris, Masson, 499 p.
- Muller, D., Bocquier, G., Nahon, D., and Paquet, H., 1981, Analyse des différenciations minéralogiques et structurales d'un sol ferrallitique à horizon nodulaire du Congo: *Cahier Office Recherche Sci. Tech. Outre-Mer (ORSTOM)*, Paris, sér. Pédologie, v. 18, 2, p. 87-109.
- Nahon, D., 1976, Cuirasses ferrugineuses et encroûtements calcaires au Sénégal occidental et en Mauritanie. Systèmes évolutifs: géochimie, structure, relais et coexistence: *Sci. Géol. Mém.*, v. 44, 232 p.
- 1985, Evolution of iron crusts in tropical landscapes, in Coleman, S. N., and Dethier, D. P., eds., *Rates of chemical weathering of rocks and minerals*: New York, Academic Press (in press).
- Nahon, D., Carozzi, A. V., and Parron, C., 1980, Lateritic weathering as a mechanism for the generation of ferruginous ooids: *Jour. Sed. Petrology*, v. 50, 4, p. 1287-1298.
- Nahon, D., Janot, C., Karpoff, A. M., Paquet, H., and Tardy, Y., 1977, Mineralogy, petrography and structures of iron crusts (ferricretes) developed on sandstones in the western part of Senegal: *Geoderma*, v. 19, p. 263-277.
- Nahon, D., Janot, C., Paquet, H., Parron, C., and Millot, G., 1979, Epigénie du quartz et de la kaolinite dans les accumulations ferrugineuses superficielles. La signification des goethites et hématites alumineuses: *Sci. Géol. Bull.*, v. 32, 4, p. 165-180.
- Nahon, D., and Millot, G., 1977, Enfoncement géochimique des cuirasses ferrugineuses par épigénie du manteau d'altération. Influence sur le paysage: *Sci. Géol. Bull.*, v. 30, 4, p. 275-282.
- Naumov, C. B., Ryzhenko, B. N., and Khodakovsky, I., 1971, *Atlas des données thermodynamiques* (en russe): Moscou, Atomizdat., 239 p.
- Nia, R., 1968, Geologische, petrographische, geochemische, untersuchungen zum problem der boehmit-diaspore-genese in guechischen oberkreide-bauxiten der Parnass-Ziona zone: Thèse, Univ. Hambourg, RFA, 133 p.
- Norrish, K., and Taylor, R. M., 1961, The isomorphous replacement of iron by aluminum in soil goethites: *Soil Sci. Jour.*, v. 12, 2, p. 294-306.
- Parks, G. A., 1972, Free energies of formation and aqueous solubilities of aluminum hydroxides and oxide hydroxides at 25°C: *Am. Mineralogist*, v. 57, p. 1163-1189.
- Pédro, G., and Delmas, A. B., 1980, Regards actuels sur les phénomènes d'altération hydrolitique: *Cahier Office Recherche Sci. Tech. Outre-Mer (ORSTOM)*, Paris, sér. Pédologie, v. 18, 3-4, p. 217-234.
- Périnet, G., and Lafont, G., 1972a, Sur les paramètres cristallographiques des hématites alumineuses: *Acad. Sci. Paris Comptes rendus*, v. 275 C, p. 1021-1024.
- 1972b, Sur la présence d'hématite alumineuse désordonnée dans les bauxites du Var. *Acad. Sci. Paris Comptes rendus*, v. 274, p. 272-274.

- Perret, D., 1983, Calcul d'équilibres géochimiques par minimisation d'énergie libre. Thèse, Ing. Doct. ENSA et Univ. Rennes, France, 120 p.
- Perret, D., Fritz, B., and Tardy, Y., 1984, MINEQ a computer program for calculating equilibria by minimization energy: *Sci. Géol. Bull.*, in press.
- Petit, J. C., Backer, L., and Herzog, E., 1964, Substitutions dans les hydroxides FeOOH, AlOOH: *Acad. Sci. Paris Comptes rendus*, v. 258, p. 4993-4994.
- Pierce, C., 1953, Computation of pore sizes from physical adsorption data: *Jour. Phys. Chemistry*, v. 57, p. 149-152.
- Pouillard, E., 1950, The behavior of alumina and of titanium oxide with oxides in iron: *Annales Chimie*, v. 5, p. 164-214.
- Rengasamy, P., Krishna-Murti, G. S. R., and Sarna, V. A. K., 1975, Isomorphous substitution of iron for aluminum in soil kaolinites: *Clays and Clay Minerals*, v. 23, p. 211-214.
- Robie, R. A., Hemingway, B. S., and Fisher, J. R., 1978, Thermodynamic properties of minerals and related substances at 298°K and one bar (10<sup>5</sup> Pascals) pressure and at higher temperatures: *U.S. Geol. Survey Bull.* 1452, 456 p.
- Robie, R. A., and Waldbaum, D. R., 1968, Thermodynamic properties of minerals and related substances at 298.15 K (25.0°C) and one atmosphere (1013 bars) pressure and at higher temperatures: *U.S. Geol. Survey Bull.* 1259, 256 p.
- Schmalz, R. F., 1959, A note on the system Fe<sub>2</sub>O<sub>3</sub>-H<sub>2</sub>O: *Jour. Geophys. Research*, v. 64, 5, p. 575-579.
- , 1968, Formation of red beds in modern and ancient deserts; discussion: *Geol. Soc. America Bull.*, v. 79, p. 277-280.
- Schwertmann, U., 1965, Zur Goethit und Hämatitbildung aus amorphen Eisen (III), hydroxyde: *Zeitschr. Pflanzenernährung, Düngung Bodenkunde*, v. 108, p. 37-45.
- , 1969, Die Bildung von Eisenoxidmineralen: *Fortschr. Mineralogie*, v. 46, p. 274-285.
- Schwertmann, U., Fischer, W. R., and Taylor, R. M., 1974, New aspects of iron oxide formation in soils: *Internat. Cong. Soil Sci.*, 11th, Moscow, Trans., p. 237-247.
- Schwertmann, U., Fitzpatrick, R. W., and Le Roux, J., 1977, Al-substitution and differential disorder in soil hematites: *Clays and Clay Minerals*, v. 25, p. 373-374.
- Schwertmann, U., and Taylor, R. M., 1977, Iron oxides, in Dixon, J. B., and Weed, S. B., eds., *Minerals in soil environments*: *Soil Sci. Soc. America*, p. 145-180.
- Shirane, G., Cox, G. E., and Roby, S. L., 1962, Mössbauer study of isomer shift, quadrupole interaction and hyperfine field in several oxides containing Fe<sup>3+</sup>: *Phys. Rev. U.S.A.*, v. 125, p. 1158-1165.
- Simon, A., Lang, M., and Seidel, M., 1959, Substitutions dans les hydroxydes FeOOH et AlOOH: *Chem. Zvesti*, v. 13, p. 733-736.
- Solymar, K., 1970, L'aluminogöethite dans les bauxites de Hongrie. Détermination de la teneur en Al incorporé au réseau: *Am. Inst. Geol. Pub. Hungary*, v. 3, p. 359-373.
- Sposito, G., 1981, *The thermodynamics of soil solutions*: Oxford, Univ. Press, 223 p.
- Tardy, Y., 1969, Géochimie des altérations. Etude des arènes et des eaux de quelques massifs cristallins d'Europe et d'Afrique: *Mém. Service Carte Géol. Alsace-Lorraine*, v. 31, 190 p.
- , 1971, Characterization of the principal weathering types by the geochemistry of waters from some European and African crystalline massifs: *Chem. Geology*, v. 7, p. 253-271.
- , 1982, Kaolinite and smectite stability in weathering conditions: *Estudios Geol.*, v. 38, 3-4, p. 295-312.
- Tardy, Y., and Fritz, B., 1981, An ideal solid solution model for calculating solubility of clay minerals: *Clay Minerals*, v. 16, p. 361-373.
- Tardy, Y., and Monnin, C., 1983, Recherches sur les mécanismes du concrétionnement. Coefficients d'activité des ions, solubilité des sels et de la silice dans les pores de petite taille: *Bull. Minéralogie*, v. 106, p. 321-328.
- Taylor, R. M., and Graley, A. M., 1967, The influence of ionic environments on the nature of iron oxides in soils: *Soil Sci., Jour.*, v. 18, p. 341-348.
- Thiel, R., 1963, Zum system  $\alpha$ -FeOOH- $\alpha$ -AlOOH: *Zeitschr. Anorg. Allg. Chemie*, v. 326, p. 70-78.
- Torrence-Martin, R., 1958, Water vapor sorption on kaolinite hysteresis: *Natl. Conf. on Clays and Clay Minerals*, 6th, p. 259-278.
- Torrent, J., Gunzman, R., and Parra, M. A., 1982, Influence of relative humidity on the crystallization of Fe (III) oxides from ferrihydrite: *Clays and Clay Minerals*, v. 30, p. 337-340.

- Ulbrich, H. H., and Merino, E., 1974, An examination of standard enthalpies of formation of selected minerals in the system  $\text{SiO}_2\text{-Al}_2\text{O}_3\text{-Na}_2\text{O-K}_2\text{O-H}_2\text{O}$ : *Am. Jour. Sci.*, v. 274, p. 610-642.
- Yvon, J., Lietard, O., Cases, J. M., and Delon, J. F., 1981, Minéralogie des argiles kaoliniques des Charentes. *Bull. Minéralogie, Supp.*, v. 104, 5, p. 55.
- Wefers, K., 1967, Phase relations in the system  $\text{Al}_2\text{O}_3\text{-Fe}_2\text{O}_3\text{-H}_2\text{O}$ : *Zeitschr. Erz. Metall.*, v. 2, p. 71-75.
- Wollast, R., 1971, Kinetic aspects of the nucleation and growth of calcite from aqueous solutions, in Bricker, O., ed., *Carbonate Cements: John Hopkins Univ. Studies in Geology*, no. 19, p. 264-273.
- Yapp, C. J., 1983, Effects of  $\text{AlOOH-FeOOH}$  solid solution on goethite-hematite equilibrium: *Clays and Clay Minerals*, v. 31, 3, p. 239-240.