

American Journal of Science

SUMMER 1994

PHANEROZOIC TRANSGRESSIONS AND REGRESSIONS ON THE CONTINENTS: A QUANTITATIVE APPROACH BASED ON AREAS FLOODED BY THE SEA AND AREAS OF MARINE AND CONTINENTAL DEPOSITION

ALEXANDER B. RONO

V. I. Vernadsky Institute of Geochemistry,
Russian Academy of Sciences, Moscow, Russia

ABSTRACT. Analysis of the areas of major tectonic units of the continents flooded by the sea during the Phanerozoic shows that there have been three major cycles of transgression and regression corresponding to the Caledonian, Hercynian, and Alpine orogenic cycles. Approximately equal areas of the platforms and geosynclinal regions have been flooded synchronously, but seas occupied only small areas of the orogenic belts. The areas of the Laurasian continental blocks flooded were more extensive than those of the Gondwanan blocks, even though the latter have an overall greater area. The flooding histories of individual cratons show notable exceptions to the global history of transgressions and regressions.

In terms of proportions of the area flooded, the platform regions average about 20 percent, the geosynclinal regions about 80 percent, and the orogenic belts about 20 percent in the Paleozoic, and 10 percent in the Mesozoic and Cenozoic. There has been a decline in the proportion of the area of marine versus continental sedimentation throughout the Phanerozoic on the continental platforms and in orogenic belts. A similar decline has occurred in the geosynclinal regions but only during the later Mesozoic and Cenozoic.

The overall pattern of flooding of the continents during the Cenozoic corresponds well with the general trend of the eustatic curve of Vail, Mitchum, and Thompson (1977). Continental uplift and subsidence calculated from the volumes of sediments eroded and deposited also correlate with the changing areas of the continents flooded by transgressing and regressing seas.

INTRODUCTION

This paper presents estimates of the areas of the platforms, geosynclines, and orogenic zones as well as the total areas of the continental blocks covered by the sea during transgressions and regressions. The estimates cover all the continents except Antarctica and all the geological record from Early Cambrian through Pliocene. Also included are estimates of the areas of marine and continental sedimentation at each of the major stages of geological evolution. Finally, the chronology of marine

transgressions and regressions is compared to vertical motions of the continental crust and changes in eustatic sealevel.

The estimates have been made by measuring the marine and land areas on the continents and volumetric calculations of the major types of sedimentary rocks shown on the series of global lithological-paleogeographical maps compiled by Ronov, Khain, and Seslavinsky (1984) for the Paleozoic and by Ronov, Khain, and Balukhovskiy (1989) for the Mesozoic and Cenozoic. The maps were compiled from literature and other data sources for regional geological studies of structure, stratigraphy, lithology, paleogeography, and paleoenvironments cited by Ronov, Khain, and Seslavinsky (1984) and by Ronov, Khain, and Balukhovskiy (1989). The paleogeography, shorelines, elevation of the land areas, and depths of water are based on interpretation of the surviving geologic record exposed in outcrop and preserved in the subsurface. Ronov, Khain, and Seslavinsky (1984) and Ronov, Khain, and Balukhovskiy (1989) took into account whether the strata were deposited under marine or continental conditions and estimated the relative elevation of the areas of erosion from the grain-size of the sediments derived from them. The original data are based on descriptions of many thousands of outcrop and borehole sections. The maps cover 28 stratigraphic intervals covering each of the epochs of the Phanerozoic. The reliability of the maps is not the same in all regions. It decreases with geologic age. The reliability also decreases from platforms to geosynclines and from the northern to the southern hemisphere within each map.

The continents are divided into three major categories of tectonic unit: platforms, geosynclines, and orogenic zones. Table 1² shows the areas of regions of erosion, regions of sedimentation, areas covered by seas, and areas covered by seas as a percentage of the total area for each category of tectonic unit. The analysis excludes the Antarctic and a few other areas that are poorly known. It should be noted that the Quaternary is not included in table 1, and that the area of continents shown as flooded on the Pliocene map of Ronov, Khain, and Balukhovskiy (1989) is much less than the area flooded at present because of the lower sealevel at the time chosen for the paleogeographic reconstruction.

FLOODING OF THE CONTINENTS

As shown in figure 1, the data demonstrate regular changes of area covered by the sea (from expansion to abrupt reduction) in the course of major tectonic cycles. The major geocratic and thalassocratic episodes in the geological history of the Earth recognized by Ronov (1968) are evident.

At present, as during the times of sealevel high stand of the Quaternary, the world ocean, including epicontinental and marginal seas, covers 361×10^6 km², corresponding to 71 percent of the Earth's surface. Seen from the perspective of geological history, this area is minimal. The

contemporary epoch should be identified as geocratic, characterized by regression from elevated continents. Ancient geocratic epochs correspond to the final stages of tectonic cycles: Early Cambrian, which corresponds to the boundary between the Baikalian and Caledonian Cycles; Late Silurian-Early Devonian, which corresponds to the boundary between the Caledonian and Hercynian Cycles; the Late Permian-Middle Triassic, which corresponds to the boundary between the Hercynian and Alpine Cycles. The Pliocene-Quaternary falls at the boundary between the Alpine and a future cycle. Figure 1 shows the successive increase in amplitude of regressions during the Phanerozoic to the extreme geocratic condition represented by the present. The middle stages of the tectonic cycles correspond to thalassocratic episodes of earth history. The areas of the continents flooded by marine transgressions were increased as a result of subsidence of the continents or shallowing of the ocean floor. The thalassocratic epochs of the Phanerozoic are the Middle Ordovician, the Middle Devonian, and the Late Cretaceous. The areas flooded decrease from the Caledonian to the Alpine cycle. Thus one common trend in the areal measurements is that the areas of the continents covered by the sea has successively decreased from the Early Paleozoic to Present. The reduction of marine area has been about $40 \times 10^6 \text{ km}^2$.

Figure 2 shows the areas of the major categories of tectonic units of the continents flooded by the sea at different times. The largest areas flooded are on the platforms, and the next largest are the geosynclinal regions. The transgressions and regressions affecting these areas correspond to the overall continental flooding episodes shown in figure 1. A second common trend of the areal measurements is that the transgressions and regressions on platforms and in geosynclinal regions are synchronous. Figure 2 also shows that the smallest areas flooded are in the orogenic regions, and here transgressions and regressions do not repeat the pattern of figure 1.

Table 2 and figure 3 summarize the data in terms of the Laurasian (Russian, Siberian, Chinese, and North American) and Gondwanan (African, Arabian, South American, Australian, Indian) platforms. Transgressions and regressions on both groups of platforms are synchronous. The synchronism is very close for the Hercynian and Alpine Cycles and differs only slightly in the Caledonian cycle, when a regression on the Gondwanan platforms corresponds to the time of maximum transgression on the Laurasian platforms. There is, however, the marked difference that the areas covered by the seas on the Laurasian platforms are much more extensive than those on the Gondwanan platforms. This relation is consistent even though the total area of Gondwanan platforms is larger than that of the Laurasian platforms.

Figure 4 compares the areas of flooding on individual platforms. As has been noted by Yanshin (1973) there is considerable asynchronicity in the development of transgressions and regressions on the different

TABLE 1
Transgressions and regressions of seas on continents and their main tectonic units during the Phanerozoic

STRATIGRAPHIC INTERVAL	DURATION OF INTERVAL (MY)	MAIN TECTONIC UNITS OF CONTINENTS	REGIONS OF EROSION (10^6 km^2)	REGIONS OF SEDIMENTATION (10^6 km^2)	AREAS COVERED BY SEAS (10^6 km^2)	AREAS COVERED BY SEAS AS % OF THE TOTAL AREA OF TECTONIC UNIT
PLIOCENE (N_2)	5.3 - 1.6 = 3.7	PLATFORMS	73.40	21.44	4.45	5
		GEOSYNCLINES	1.32	1.58	0.99	34
		OROGENIC BELTS	21.23	14.23	1.91	5
		WHOLE CONTINENT	95.95	37.25	7.35	5
MIOCENE (N_1)	23.7 - 5.3 = 18.4	PLATFORMS	74.55	20.14	6.53	7
		GEOSYNCLINES	0.74	2.13	2.10	73
		OROGENIC BELTS	20.47	14.64	4.30	12
		WHOLE CONTINENT	95.76	36.91	12.93	10
OLIGOCENE (P_3)	36.6 - 23.7 = 12.9	PLATFORMS	77.80	17.71	5.85	6
		GEOSYNCLINES	1.76	3.91	3.58	63
		OROGENIC BELTS	21.26	9.23	2.76	9
		WHOLE CONTINENT	100.82	30.85	12.19	9
EOCENE (P_2)	57.8 - 36.6 = 21.2	PLATFORMS	75.35	24.31	14.67	15
		GEOSYNCLINES	2.27	7.95	7.18	70
		OROGENIC BELTS	14.73	8.00	2.19	10
		WHOLE CONTINENT	92.35	40.26	24.04	18
PALEOCENE (P_1)	66.4 - 57.8 = 8.8	PLATFORMS	77.37	20.32	12.35	13
		GEOSYNCLINES	3.82	6.13	6.03	61
		OROGENIC BELTS	19.06	5.86	1.87	7
		WHOLE CONTINENT	100.20	32.31	20.25	15
LATE CRETACEOUS (K_2)	97.5 - 66.4 = 31.1	PLATFORMS	56.90	41.76	22.41	23
		GEOSYNCLINES	1.37	16.82	14.53	80
		OROGENIC BELTS	9.28	7.45	1.81	11
		WHOLE CONTINENT	67.55	66.03	38.72	29
EARLY CRETACEOUS (K_1)	144 - 97.5 = 46.3	PLATFORMS	59.77	40.06	17.04	17
		GEOSYNCLINES	3.55	18.76	15.12	70
		OROGENIC BELTS	9.59	7.06	0.78	5
		WHOLE CONTINENT	72.91	65.88	33.54	24

LATE JURASSIC (J ₃)	163 - 144 = 19	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	74.30 4.91 8.13 87.34	25.15 16.05 4.12 45.32	16.82 14.10 1.44 32.36	17 67 12 24
MIDDLE JURASSIC (J ₂)	187 - 163 = 24	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	75.08 2.29 11.54 88.91	25.47 12.76 6.04 44.27	14.83 12.49 1.40 28.79	15 83 8 22
EARLY JURASSIC (J ₁)	208 - 187 = 21	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	79.85 1.54 10.85 92.24	22.97 10.93 5.73 39.63	11.40 10.86 1.35 23.61	11 87 8 18
LATE TRIASSIC (T ₃)	230 - 208 = 22	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	78.75 1.56 9.81 90.12	23.53 13.63 3.63 40.79	10.44 13.27 0.50 24.21	9 87 4 18
MIDDLE TRIASSIC (T ₂)	240 - 230 = 10	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	82.01 1.69 23.19 96.89	17.95 11.57 1.77 31.29	10.65 11.36 0.69 22.70	11 86 5 18
EARLY TRIASSIC (T ₁)	245 - 240 = 5	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	79.57 1.94 10.52 92.03	24.73 10.22 1.65 36.60	11.49 9.96 0.07 21.52	11 82 1 17
LATE PERMIAN (P ₂)	258 - 245 = 13	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	75.61 1.07 14.58 91.26	19.34 9.23 4.88 33.45	11.55 8.94 1.81 22.30	12 86 9 18
EARLY PERMIAN (P ₁)	286 - 258 = 28	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	68.88 0.98 13.07 82.93	24.70 9.36 7.73 41.79	13.96 9.18 4.02 27.16	15 89 19 22

TABLE 1
(continued)

STRATIGRAPHIC INTERVAL	DURATION OF INTERVAL (MY)	MAIN TECTONIC UNITS OF CONTINENTS	REGIONS OF EROSION (10^6 km^2)	REGIONS OF SEDIMENTATION (10^6 km^2)	AREAS COVERED BY SEAS (10^6 km^2)	AREAS COVERED BY SEAS AS % OF THE TOTAL AREA OF TECTONIC UNIT
LATE & MIDDLE CARBONIFEROUS (C_{3+2})	320 - 286 = 34	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	68.26 1.02 11.53 80.81	28.04 9.66 8.81 46.51	16.69 9.46 5.36 31.51	17 89 26 25
EARLY CARBONIFEROUS (MISSISSIPPIAN)	360 - 320 = 40	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	71.91 2.38 8.67 82.96	20.99 17.63 4.03 42.65	15.92 17.28 2.36 35.46	17 86 19 28
LATE DEVONIAN (D_3)	374 - 360 = 14	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	67.65 2.50 7.32 77.47	25.39 18.55 4.10 48.04	21.80 17.99 2.23 42.02	23 85 19 33
MIDDLE DEVONIAN (D_2)	387 - 374 = 13	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	68.75 1.83 7.49 78.17	26.17 18.82 3.57 48.56	24.23 18.43 2.04 44.70	25 89 18 35
EARLY DEVONIAN (D_1)	408 - 387 = 21	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	74.53 2.39 6.94 83.86	16.90 17.48 4.18 38.56	14.51 17.29 2.45 34.25	16 87 22 28
LATE SILURIAN (S_2)	421 - 408 = 13	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	72.74 2.89 8.16 83.79	22.29 15.24 2.12 39.65	17.65 14.93 1.65 34.23	19 82 16 28
EARLY SILURIAN (S_1)	438 - 421 = 17	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	66.75 2.92 4.70 74.37	28.38 17.08 2.13 47.59	27.56 16.82 1.71 46.09	29 84 25 38

LATE ORDOVICIAN (O ₃)	458 - 438 = 20	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	69.49 3.84 2.14 75.47	26.85 18.52 0.33 45.70	22.50 18.24 0.27 41.01	23 82 11 34
MIDDLE ORDOVICIAN (O ₂)	478 - 458 = 20	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	69.28 2.46 1.40 73.14	27.90 19.78 0.41 48.09	27.44 19.51 0.26 47.21	26 88 14 39
EARLY ORDOVICIAN (O ₁)	505 - 478 = 27	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	70.26 2.38 1.21 73.85	25.40 18.32 0.46 44.18	24.65 18.32 0.38 43.35	26 88 23 37
* LATE CAMBRIAN (€ ₃)	520 - 505 = 15	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	66.54 2.67 6.02 75.23	24.43 15.61 0.92 40.96	22.93 15.46 0.46 38.85	25 84 7 33
MIDDLE CAMBRIAN (€ ₂)	540 - 520 = 20	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	66.77 2.01 6.76 75.54	25.65 14.47 1.03 41.15	22.17 14.41 0.45 37.03	24 87 6 32
EARLY CAMBRIAN (€ ₁)	570 - 540 = 30	PLATFORMS GEOSYNCLINES OROGENIC BELTS WHOLE CONTINENT	55.78 1.78 15.87 73.43	22.94 13.96 1.76 38.66	18.23 13.86 1.28 33.37	23 88 7 30


Areas of continents covered by seas, 10^6 km^2 

Fig. 1. Transgressions and regressions of the seas on the continents during the Phanerozoic.

platforms. The synchronicity of transgressions and regressions on platforms and in geosynclinal regions apparent in the global data suggests that there are global rhythms that result from periodic increase and decline of tectonic activity. The global rhythms dominate the evolution of


Areas of continents covered by seas, 10^6 km^2 

Fig. 2. Transgressions and regressions of the seas over platforms, geosynclines, and orogenic belts during the Phanerozoic.

vast lithospheric regions (Ronov, Khain, and Balukhovskiy, 1976), such as Laurasia and Gondwana, as shown in figure 3. The global rhythms are apparent in the supercontinental aggregations, although there is a much more noisy signal in the data from individual platforms.

TABLE 2
Transgressions and regressions of the seas on Laurasian and Gondwanan platforms during the Phanerozoic

STRATIGRAPHIC INTERVAL	DURATION OF INTERVAL (MY)	GROUPS OF PLATFORMS	REGIONS OF EROSION (10^6 km^2)	REGIONS OF SEDIMENTATION (10^6 km^2)	AREAS COVERED BY SEAS (10^6 km^2)	AREAS COVERED BY SEAS AS % OF THE TOTAL AREA OF TECTONIC UNIT
PLIOCENE (N_2)	5.3 - 1.6 = 3.7	LAURASIAN	31.80	8.63	2.69	7
		GONDWANAN	41.60	12.81	1.76	3
MIOCENE (N_1)	23.7 - 5.3 = 18.4	LAURASIAN	32.14	7.93	3.79	9
		GONDWANAN	42.41	12.21	2.74	5
OLIGOCENE (P_1)	36.6 - 23.7 = 12.9	LAURASIAN	30.92	9.20	4.42	11
		GONDWANAN	46.88	8.51	1.43	3
Eocene (P_2)	57.8 - 36.6 = 21.2	LAURASIAN	32.51	11.93	9.42	21
		GONDWANAN	42.84	12.38	5.25	9
PALEOCENE (P_1)	66.4 - 57.8 = 8.8	LAURASIAN	32.22	9.89	7.81	18
		GONDWANAN	45.10	10.43	4.54	8
LATE CRETACEOUS (K_2)	97.5 - 66.4 = 31.1	LAURASIAN	23.16	31.64	14.04	26
		GONDWANAN	33.74	10.12	8.37	19
EARLY CRETACEOUS (K_1)	144 - 97.5 = 46.3	LAURASIAN	28.65	19.52	10.91	23
		GONDWANAN	31.13	20.54	6.13	12
LATE JURASSIC (J_3)	163 - 144 = 19	LAURASIAN	27.35	16.25	13.17	30
		GONDWANAN	46.95	8.90	3.65	6
MIDDLE JURASSIC (J_2)	187 - 163 = 24	LAURASIAN	27.17	17.48	11.11	25
		GONDWANAN	47.91	7.99	3.72	7
EARLY JURASSIC (J_1)	208 - 187 = 21	LAURASIAN	31.48	15.58	8.29	18
		GONDWANAN	48.37	7.39	3.11	6
LATE TRIASSIC (T_3)	230 - 208 = 22	LAURASIAN	30.71	16.06	8.24	18
		GONDWANAN	48.04	7.47	2.20	4
MIDDLE TRIASSIC (T_2)	240 - 230 = 10	LAURASIAN	33.87	12.50	9.17	20
		GONDWANAN	48.14	5.45	1.48	3
EARLY TRIASSIC (T_1)	245 - 240 = 5	LAURASIAN	31.43	18.30	10.60	21
		GONDWANAN	48.14	6.43	0.89	2

LATE PERMIAN	(P ₂)	258 - 245 = 13	LAURASIAN	30.21	13.12	9.03	21
EARLY PERMIAN	(P ₁)	286 - 258 = 28	GONDWANAN	45.40	6.22	2.52	5
PERMIAN			LAURASIAN	18.37	13.73	10.47	33
LATE & MIDDLE (C ₃₊₂)			GONDWANAN	40.51	10.97	3.49	7
CARBONIFEROUS		320 - 286 = 34	LAURASIAN	27.03	14.42	10.86	26
EARLY CARBONIFEROUS	(C ₁)		GONDWANAN	41.23	13.58	5.83	11
			LAURASIAN	26.05	13.56	11.53	29
CARBONIFEROUS		360 - 320 = 40	GONDWANAN	45.86	7.43	4.39	8
LATE DEVONIAN	(D ₃)	374 - 360 = 14	LAURASIAN	24.96	15.85	14.76	36
MIDDLE DEVONIAN	(D ₂)		GONDWANAN	42.69	9.54	7.04	13
			LAURASIAN	26.96	14.66	13.48	32
DEVONIAN		387 - 374 = 13	GONDWANAN	41.79	11.51	10.75	20
EARLY DEVONIAN	(D ₁)		LAURASIAN	31.99	7.73	6.65	17
		408 - 387 = 21	GONDWANAN	42.54	9.17	7.86	15
LATE SILURIAN	(S ₂)	421 - 408 = 13	LAURASIAN	30.85	10.71	9.35	22
EARLY SILURIAN	(S ₁)		GONDWANAN	41.89	11.58	8.30	15
			LAURASIAN	26.04	16.26	15.81	37
SILURIAN		438 - 421 = 17	GONDWANAN	40.71	12.12	11.75	22
LATE ORDOVICIAN	(O ₃)	458 - 438 = 20	LAURASIAN	25.09	18.35	17.9	41
			GONDWANAN	44.40	8.50	4.60	9
MIDDLE ORDOVICIAN	(O ₂)	478 - 458 = 20	LAURASIAN	24.89	20.06	19.90	44
EARLY ORDOVICIAN	(O ₁)		GONDWANAN	44.39	7.84	7.54	14
			LAURASIAN	24.74	18.41	18.41	43
ORDOVICIAN		505 - 478 = 27	GONDWANAN	45.52	6.99	6.24	12
LATE CAMBRIAN	(C ₂)	520 - 505 = 15	LAURASIAN	24.53	18.27	17.36	41
			GONDWANAN	42.01	6.16	5.57	12
MIDDLE CAMBRIAN	(C ₁)	540 - 520 = 20	LAURASIAN	22.73	19.90	17.54	41
			GONDWANAN	44.04	5.75	4.63	6
EARLY CAMBRIAN		570 - 540 = 30	LAURASIAN	22.00	17.37	14.56	37
			GONDWANAN	33.29	5.57	3.67	9


Areas of platforms covered by seas, 10^6 km^2 

Fig. 3. Transgressions and regressions of the seas on Laurasian and Gondwanan platforms during the Phanerozoic.

PROPORTIONS OF THE CONTINENTS FLOODED

Figure 5 shows the proportion of the total area of the continents covered by the sea. The flooded areas of the continents range between 3 and 40 percent of the total continental area. Most of the continental blocks have been land throughout the entire Phanerozoic. The general


Areas of platforms covered by seas, 10^6 km^2 

Fig. 4. Transgressions and regressions of the seas on the major cratonic platforms during the Phanerozoic.

trend toward increase of land area on the continental blocks through the Phanerozoic is again apparent.

The stable (platform) and mobile (geosynclines and orogenic belts) regions of the continents have had very different land/sea distributions. The land area greatly predominates on the platforms, as shown in figure

Percent of total area of continents




Fig. 5. Distribution of land (including areas of continental sedimentation) and sea as a percentage of the total area of the continents during the Phanerozoic.

6. On the platforms the marine areas generally did not exceed a quarter of the total area; the flooding ranged from 5 to 30 percent. In contrast, the geosynclines are characterized by a dominance of marine conditions. Paleozoic seas covered up to 90 percent of the total area of the geosyn-

Percent of total area of platforms




Fig. 6. Distribution of land (including areas of continental sedimentation) and sea as a percentage of the area of the continental platforms during the Phanerozoic.

clines, as shown in figure 7. Only in the Late Mesozoic and the Cenozoic did the marine area of the geosynclinal regions decrease significantly. The largest regressions were in the Early Cretaceous and Pliocene. As shown in figure 8, the marine areas in the orogenic belts were very

Percent of total area of geosynclines




Fig. 7. Distribution of land (including areas of continental sedimentation) and sea as a percentage of the area of geosynclines during the Phanerozoic.

restricted relative to the platforms and geosynclinal regions. The areas covered by the sea within orogenic belts were more extensive during the Paleozoic than during the Mesozoic or Cenozoic. The geosynclines, which are the regions of greatest rate and magnitude of crustal subsi-

Percent of total area of orogenic zones




Fig. 8. Distribution of land (including areas of continental sedimentation) and sea as a percentage of the area of orogenic belts during the Phanerozoic.

dence (Ronov, 1980, in press; Ronov, Khain, and Balukhovskiy, 1976; Ronov, Yaroshevskiy, and Migdisov, 1990) are predominantly marine and serve as the route by which waters from the world ocean reach the continental platforms.

RATIOS OF MARINE AND CONTINENTAL SEDIMENTATION

Comparison of the areas of marine and continental sedimentation on the continents shows a long-term trend for reduction of the area of marine sedimentation and increase of the area of continental sedimentation during the Phanerozoic, as shown in figure 9. This trend is best developed on the platform areas, as shown in figure 10. It is poorly developed in the geosynclinal regions, as shown in figure 11. The geosynclinal regions are predominantly marine, and appreciable areas of continental sedimentation do not appear until the middle Mesozoic. The orogenic belts do show the trend toward replacement of areas of marine sedimentation by areas of continental sedimentation, as shown in figure 12. The shift from marine to continental sedimentation is especially apparent in the Alpine Cycle.

The measurements of areas flooded can be interpreted in terms of transgressions and regressions, corresponding broadly to the Caledonian, Hercynian, and Alpine tectonic cycles. The curve of areas of continents covered by the sea is compared with the eustatic sealevel curve of Vail, Mitchum, and Thompson (1977) in figure 13. Comparison of the two curves indicates that the flooding of the continents generally corresponds to the rises in sealevel and regression from the continents corresponds to falling sealevel of Vail, Mitchum, and Thompson (1977). The coincidence of the two curves is evidence that the transgressions and regressions recorded in changes of areas of the continents flooded is a reflection of eustatic sealevel changes. The data on which the two curves are based are independent: the curve of area of the continents flooded is based on measurements of the maps of Ronov, Khain, and Seslavinsky (1984) and Ronov, Khain, and Balukhovskiy (1989), and the sealevel curve of Vail, Mitchum, and Thompson (1977) is based on seismic stratigraphic studies of continental margins.

FLOODING, UPLIFT, AND SUBSIDENCE

Figure 14 shows the changes in areas of the seas on the continents and apparent subsidence or elevation of the continental blocks. Ronov (1980, table 13) gave the results of calculating the amounts of uplift of the areas of continents undergoing erosion that would be required to yield a volume of eroded material equal to that deposited in contemporaneous continental and marine sedimentary basins. The total average amounts of uplift of the areas undergoing erosion for each stratigraphic interval is shown in figure 14. Similarly, the average subsidence of the continental areas in which sedimentation occurred was calculated by Ronov (1980, table 13). The total average amounts of subsidence for each of the stratigraphic intervals is shown in figure 14.

Subsidence of the continents or elevation of the ocean floor results in transgression and flooding of the continents. Elevation of the continents or subsidence of the ocean floor results in regression and reduction of the

Total area of sedimentation zones of continents




Fig. 9. Change in the ratio of areas of marine and continental sedimentation over the whole continents during the Phanerozoic.

area covered by the sea. The area flooded also depends on the amount of ice stored in polar ice sheets during glaciations. Figure 14 shows that the maximal regressions occurred during the Pliocene, the Late Permian, and the Late Ordovician and may correspond to times of glaciation.




Fig. 10. Change in the ratio of areas of marine and continental sedimentation on the continental platforms during the Phanerozoic.

CAUSES OF EUSTATIC SEALEVEL CHANGE

Donovan and Jones (1979) suggested that the oscillations of sealevel might be connected with changes in the morphology of the ocean floor. They noted that the volume of the ocean basins would change in

Total area of geosynclinal sedimentation




Fig. 11. Change in the ratio of areas of marine and continental sedimentation in geosynclines during the Phanerozoic.

response to evolution and changes in the volume of the mid-ocean ridge system. Unfortunately, knowledge of the mid-ocean ridge system is available only for Late Jurassic and younger epochs, and the relation of older sealevel changes to changes in the volume of the ocean basins must,

Total area of orogenic sedimentation




Fig. 12. Change in the ratio of areas of marine and continental sedimentation in orogenic belts during the Phanerozoic.

because of the lack of data, remain hypothetical. The problem of what drives the transgressions and regressions of the sea is far from a final solution. It is hoped that the data presented here will stimulate thought toward finding a solution of this fundamental problem of geology.




Fig. 13. Transgressions and regressions of seas on the continents (Ronov, in press) and changes in sea level during the Phanerozoic (Vail, Mitchum, and Thompson, 1977).


Areas of continents covered by seas, 10^6 km^2 

Fig. 14. Transgressions and regressions of seas, average uplift of areas of erosion, and average subsidence of areas of sedimentation on the continents. Subsidence was calculated by dividing the volumes of rock deposited by the area over which the rocks were deposited. Uplift was calculated by dividing the volumes of rock deposited by the land area over which erosion occurred (Ronov, 1980). Major glaciations are indicated.

ACKNOWLEDGMENTS

The author is indebted to Bryan Gregor, Dork Sahagian, and William Hay for their constructive reviews of the draft of the manuscript. He also thanks Christopher Wold and Barry Eakins for redrafting the figures. The author and the editors of the *American Journal of Science* thank William Hay for his assistance in editing the manuscript.

REFERENCES

- Donovan, D. T., and Jones, E. J. W., 1979, Causes of world-wide changes in sea level: *Journal of the Geological Society* (London), v. 136, p. 187–192.
- Ronov, A. B., 1968, Probable changes in the composition of sea water during the course of geologic time: *Sedimentology*, v. 10, p. 25–43.
- 1980, The earth's sedimentary shell (quantitative patterns of its structure, compositions, and evolution).—The 20th V. I. Vernadski Lecture, March 12, 1978 (in Russian), in Yaroshevskii, A. A., editor, *The Earth's Sedimentary Shell (Quantitative Patterns of Its Structure, Compositions, and Evolution)*: Moscow, Nauka, p. 80 (English translation in *International Geology Review*, 24, 1313–1388, 1982, also *American Geological Institute Reprint Series* 5, 73 p., 1983).
- Ronov, A. B., in press. *The stratosphere or sedimentary shell of the Earth*: Moscow, Izdatelstvo Nauka (in Russian).
- Ronov, A. B., Khain, V. E., and Balukhovskiy, A. N., 1976, Changes in the distribution of volumes and rates of deposition of sedimentary and volcanogenic deposits during Phanerozoic time (within the present continents): *Academic Nauk SSSR, Izvestia, Seriya Geologicheskaya*, n. 12, p. 5–12 (in Russian).
- 1989, *Atlas of Lithological-Paleogeographical Maps of the World: Mesozoic and Cenozoic of Continents and Oceans*: Moscow, Editorial Publishing Group, 79 p. (in Russian).
- Ronov, A. B., Khain, V. E., and Seslavinsky, K. B., 1984, *Atlas of Lithological-Paleogeographical Maps of the World: Precambrian and Paleozoic of Continents*: Leningrad, Academy of Sciences Press, 70 p. (in Russian).
- Ronov, A. B., Yaroshevskiy, A. A., and Migdisov, A. A., 1990, The chemical structure of the Earth's crust and the geochemical balance of main elements: Moscow, Izdatelstvo Nauka, 182 p. (in Russian).
- Vail, P. R., Mitchum, R. M., Jr., and Thompson, S., III, 1977, Global cycles of relative changes of sea level, in Payton, C. E., editor, *Seismic Stratigraphy—Applications to Hydrocarbon Exploration*, *American Association of Petroleum Geologists Memoir* 26, p. 83–97.
- Yanshin, A. L., 1973, On the so called global transgressions and regressions. *Biulletin Moskovskoe Obshchestvo Ispytalelei Prirody, Otdel Geologicheskii*, v. 48 (n. 2), p. 9–44 (in Russian).